

RESEARCH

Open Access



The impact of COVID-19 on the changes in health behaviours among Black, Asian and Minority Ethnic (BAME) communities in the United Kingdom (UK): a scoping review

Ifunanya Nduka¹, Isabella Kabasinguzi¹, Nasreen Ali^{1*}, Peter Ochepo¹, David Abdy¹, Erica Jane Cook², Chimeme Egbutah³, Sally Cartwright³ and Gurch Randhawa¹

Abstract

Background The COVID-19 pandemic has led to changes in health behaviours, which include eating patterns and nutrition, smoking, alcohol consumption, sleeping patterns, physical activity and sedentary behaviour. There is a dearth of evidence reporting the impact of COVID-19 on the health behaviour of Black, Asian and minority ethnic (BAME) communities. This scoping review synthesises the available evidence on the impact of COVID-19 on the changes in health behaviours among BAME communities in the UK.

Methods Following a keyword strategy, 16 electronic databases were searched for articles that met the screening criteria. These articles were then reviewed in full text. Empirical studies that assessed COVID-19 related health behaviour changes among BAME communities in the UK, conducted during the COVID-19 pandemic between July 2020 and August 2021 and published in English language, were set as inclusion criteria. An initial 2160 studies were identified in the selected databases. After removing duplications and screening the title and abstracts of the 2154 studies, only 4 studies were selected to be reviewed as they met the inclusion criteria. The included studies employed different sample sizes which ranged from $N=47$ to $N=30,375$ and reported several health behaviour changes. Out of the 4 included studies, 3 studies included BAME groups within their sample as a subgroup while one study focused specifically on BAME groups.

Results The scoping review found that there were lower levels of physical activity among BAME groups compared to the White ethnic groups. About 41.7% of BAME groups reported drinking less alcohol than usual compared to their white counterparts who were 34%. Study participants from BAME backgrounds had the greatest effect of COVID-19 on decisions to purchase healthier food compared to those from white backgrounds whose decisions on purchasing healthier food were least affected. Some participants reported an increase in positive hygiene practices due to the COVID-19 pandemic.

Conclusion COVID-19 had a significant impact on the health behaviours of BAME groups especially during the lockdowns as they reported changes to behaviour such as low levels of physical activities. Hence, it is important to promote health awareness among BAME groups to encourage healthy living. In addition, programmes such as physical

*Correspondence:

Nasreen Ali
nasreen.ali@beds.ac.uk

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

fitness activities that favour BAME groups should be put in place, for example BAME women's walking groups to encourage people from BAME backgrounds to engage in physical activities. Furthermore, healthy food programmes such as food parcels can be given to people from BAME backgrounds who are not able to afford healthy food due to the impact of COVID-19. Nonetheless, the COVID-19 pandemic has increased positive hygiene among BAME groups which is important in preventing other diseases and infections.

Keywords Covid-19, BAME, Health behaviors, Change, UK

Introduction

The COVID-19 pandemic has led to changes in health behaviours and consequently, health outcomes [1]. Health behaviours are intentional or unintentional actions or habits taken by individuals that affect health or mortality [2]. Health behaviours include physical activity, smoking, alcohol use, sleep, diet, healthcare-seeking behaviours, adherence to medical treatments and sedentary behaviours [2, 3]. Changes in health behaviour through disruptions to daily routines such as reduced physical activity and COVID-19 related anxiety impact on health and wellbeing [3, 4].

Before COVID-19, a decrease in preventative health behaviours accounted for 40% of the causes of premature mortality [4]. According to the World Health Organisation (WHO), health behaviours and lifestyle factors are a major contributory factor towards non-communicable diseases such as diabetes and cardiovascular diseases [5, 6]. In addition, epidemiological studies have reported that preventative health behaviours can delay the risk of premature mortality and morbidity [7]. Furthermore, the combined effects of four defined health behaviours (smoking, alcohol consumption, fruit and vegetable intake and physical activity) predicts a four-fold difference in the risk of dying over an eleven year time period for older and middle-aged people [8].

Available evidence argues that the COVID-19 social distancing measures including the two lockdowns on March 2020 and November 2020 [9], have affected health behaviours [10, 11] with social distancing having a substantial impact on decreased physical activity [12–14], dietary choices specifically the consumption of less healthy foods such as high sugar calorie-rich foods [15–17], increase in alcohol and tobacco consumption [18–21] and decrease in overall emotional wellbeing [22]. Sedentary behaviours such as prolonged sitting and screen time, has also increased due to social measures and constraints resulting from COVID-19 [1]. Studies have suggested that prolonged sedentary behaviours may increase hormonal dysfunction and cardiometabolic risks, which may aggravate underlying chronic conditions [23, 24]. Physical inactivity, as well as sedentary behaviours are independent

determinants for poor cardiometabolic health [25]. Sedentary behaviours such as sitting time increases the susceptibility of cardiovascular diseases and mortality [3, 10].

Negative changes in health behaviours are likely to interact in complex ways. For example, those smoking tobacco may also be likely to drink more alcohol and engage in less physical activity [26]. These interacting behaviours are very likely to be associated with existing multiple disadvantage and health inequalities, such as lower socioeconomic status [27, 28]. The literature that discussed BAME communities and health behaviours focusses on the differences in preventive health behaviours between people from BAME background and people from white British background during COVID-19 [29]. For instance, a study exploring changes during the COVID-19 lockdown reported that a higher percentage of people from BAME background than those from white background said they would not take the COVID-19 vaccine [30].

BAME groups have been disproportionately affected by COVID-19 [29]. For the purposes of this paper. We are aware that the term BAME is contested and criticised because it risks homogenising the experience of people from different ethnic backgrounds and masking inequalities between groups; it also maintains white British ethnic identity as the dominant privileged norm [31]. We recognise that BAME groups are not homogeneous, and that individuals from these communities have their own narratives [32]. This disproportionate impact on BAME groups have been attributed to an intersection of environmental, physiological, deprivational as well as health-related behaviours [33].

Comparing ethnic differences in health behaviour during COVID-19, the literature highlights that BAME groups have shown irregular sleep patterns, decreased physical activity and less consumption of fruits and vegetables [34] than white British [35]. Also, BAME groups have shown lesser alcohol consumption during COVID-19, compared to their White counterparts [35]. Even though sales figures of alcohol indicate an overall increase in the consumption of alcohol [36–38]. An explanation for the low alcohol intake noted among

people from BAME group during COVID-19, may be due to social norms among the BAME communities [39]. Evidence suggests that the differences in health behaviours may be associated with the disproportionate impact of COVID-19 on BAME groups [29]. This has caused increasing concerns on the long-term effects of negative health behaviour changes on mental and physical health outcomes [40].

There is limited evidence on the impact of COVID-19 on health behaviours among the BAME groups [41, 42]. Therefore, this scoping review aimed to synthesise the evidence on the impact of COVID-19 on the changes in health behaviours and lifestyle of BAME groups in the UK. The findings from this scoping review are important for policy makers, local councils, and commissioners to develop and commission programmes and interventions that will support the BAME communities improve health and wellbeing and tackle health inequalities.

Methods

We conducted a systematic scoping review of published peer-reviewed articles reporting the impact of COVID-19 on health behaviours among BAME communities. We followed the methodological framework outlined by Arksey and O'Malley [43] for scoping reviews to guide our response to the research question of "What are the COVID-19 related health behaviour changes among BAME communities in the UK?". The framework included identifying relevant studies, selecting studies, charting/extracting the data, collating, summarising, and reporting the results. Scoping studies are used to provide in-depth or comprehensive coverage (breadth) of available literature. Following the literature search on the COVID-19-related health behaviour changes among BAME communities, it was clear that there is insufficient evidence assessing and synthesising health behaviour changes among BAME groups. As a result, this scoping review focused on the breadth of relevant literature. Further, due to inadequate evidence in the previous search for literature, it was expected that the final included studies for this review were going to be heterogeneous, thus, the need for a scoping review.

Identifying the relevant studies

Search strategy

The search strategy was developed according to the research question and definition of key concepts. Relevant keywords were created under the categories of population (BAME group), region, COVID-19 and outcome (health behaviour changes). Search terms which were alternative words to keywords were formulated. These are presented in supplemental file 1. The search terms were combined with Boolean

operators to search for empirical studies published between July 2020 and August 2021, in electronic databases. The Librarian was consulted in formulating search strings. Relevant electronic databases were identified using the University of Bedfordshire digital library under healthcare-related databases. These databases included AMED, British Nursing database, CINAHL plus with full text, Medline, PsycINFO, Pubmed central, Cochrane library, TRIP database, UK Pubmed central, socINDEX, Annual reviews, ISI Web of Science, Academic search complete, Credo reference, Sage premier and Scopus. In addition, the search process was extended to other website platforms such as Google and Google scholar. Reference lists of included studies were also searched manually (snowballing) for more relevant papers until a saturation point was reached where no new papers were being identified.

Selecting the studies

The inclusion criteria applied included papers that assessed COVID-19 related health behaviour changes, papers that assessed BAME communities, papers written in English language, papers conducted in the UK, papers conducted during COVID-19. The exclusion criteria applied were papers that were not written in English language, papers that were conducted outside the UK, and papers that were not conducted during COVID-19.

At first, the selected studies were screened by their title and abstract. This resulted in removal of papers that did not meet the inclusion criteria. Duplicates were also removed. Finally, the relevant papers were screened by their full texts and final decisions were made by the researchers on the final papers to be included in the review. A flow diagram showing the different stages of study selection based on PRISMA, is shown in Fig. 1.

Data extraction

After the selection of the final studies, a data extraction spreadsheet based on the Cochrane data extraction was used at this stage of the scoping review. The data extraction spreadsheet was refined and developed to accommodate relevant information related to the research question of this scoping review. Thus, enabling the provision of a comprehensive and systematic extraction of information [44]. The data extraction spreadsheet included the following headings: date, authors, article number, publication type, description, methods, design, setting, outcomes (health behaviour changes) and bias identified. The data were extracted manually into the data extraction spreadsheet template




Fig. 1 PRISMA flow diagram showing the study selection process

after it was created and designed using Microsoft Word by IN. The data extraction process was reviewed by all authors while considering the scoping review research question and objectives. Disagreements were resolved through discussion and consensus and an agreement for final inclusion was reached by all authors.

Charting and summarising the findings

A descriptive-analytical narrative method [43] was used to chart the findings of this scoping review. A spread sheet was developed using Microsoft excel to systematically chart the findings across the heterogeneous studies. This charting process involved extracting information from the included studies in line with the objectives of the scoping review. Therefore, the charting process recorded the health behaviour changes of BAME groups, the authors, year of publication, study setting and methods, participants and other findings.

Results

Table 1 shows the features of the final included study articles. The study articles were all published in 2021, which could be attributed to the recent status COVID-19. The study articles included participants

ranging from $N=47$ to $N=30,375$. Three study articles were quantitative studies [45–47] while one study article was a qualitative study [48]. All study articles recruited participants and collected data through online platforms such as Zoom, Microsoft Teams and application software, due to the COVID-19 restrictions. The employed research approaches were online cross-sectional surveys [45–47], secondary longitudinal and exploratory analysis [47] and in-depth qualitative study [48]. Furthermore, 3 study articles reported changes in individual health behaviours while one study article reported generally on changes in health behaviour among Muslim communities [48]. The individual health behaviour changes that were reported are as follows; physical activity [47], alcohol drinking behaviour [45] and eating and purchasing behaviours [46].

Discussion

We synthesised the evidence on the impact of COVID-19 on the changes in health behaviours among BAME groups in the UK and we recorded the different methodologies participants and findings used in included studies. The included studies showed that people from BAME groups did make up part of the study sample in each study but only one study focused specifically on BAME groups [4] and thus there is a paucity of research addressing COVID-19 related health behaviour changes among the BAME communities in the UK.

The included studies showed both positive and negative changes in health behaviours and lifestyle patterns due to COVID-19 among BAME population.

Only one of the included studies [47] reported changes in physical activity and showed lower levels of physical activity in BAME population when compared to white population. This is not uncommon, as studies conducted in other countries have also reported negative changes in physical activity during the COVID-19 lockdowns. In a cross-sectional survey in Zimbabwe, it was reported that more than half of the participants reduced their physical activity during COVID-19 [49]. Similarly, Rodriguez-Perez *et al.* [50] highlighted that an estimated 60% of their participants reduced their levels of physical activity. This may be explained by evidence that describes an increase in screen time [49], working from home or closures and restrictions on gym centres and sporting activities during the COVID-19 lockdowns [51]. Furthermore, a survey on US students comparing data from 2018/2019 to data collected during the 2020 lockdown has shown decreased levels of physical activity showing the negative effect of COVID-19 lockdown on young people [52].

According to Garnett *et al.* [45], people from BAME groups were more likely to drink less alcohol than usual during the COVID-19 lockdowns. Some BAME groups

Table 1 The features of the final included study articles

S.no	Authors	Aim	Method	Participants (N)	Main findings
1	Wickersham et al. (2021) [47]	To describe the longitudinal trajectories of physical activity following the start of lockdown among students at a large UK university, and to investigate whether these trajectories varied according to age, gender and ethnicity	Secondary, longitudinal, and exploratory data analysis of the RMT King's Move physical activity tracker app	770 (129 = White, 7 = Black, 77 = Asian, 21 = mixed, 31 = others)	<ul style="list-style-type: none"> -Lower levels of physical activity in BAME groups -Gradual increase in the number of steps walked per week following the commencement of the UK nationwide lockdown -Decrease in the number of miles run by females with no change among males. -41.7% of BAME groups reported drinking less alcohol than usual. 34.0% of White groups reported drinking less than usual -37.1% of BAME groups reported drinking more than usual. 35.1% white groups reported drinking more than usual -BAME groups were more likely than those of white ethnicity to be drinking less than usual -Significant association between drinking and being younger, female, high income, stress about catching or becoming ill from COVID-19 and having an anxiety disorder -Psychological factors predicted changes in drinking behaviours
2	Garnett et al. (2021) [45]	To assess what factors were associated with reported changes to usual alcohol drinking behaviour during the start of lockdown in the UK	A cross-sectional analysis of baseline survey data from the University College London (UCL)	30,375	<ul style="list-style-type: none"> -BAME groups had the greatest effect of COVID-19 on decisionmaking and purchasing of healthier foods compared to participants from white backgrounds - Among the BAME groups, a considerable number of people from mixed or multiple ethnic groups had the lowest number of participants who had their decisions on healthier food affected "moderately or a little bit"
3	Ogundijo, D. A., Tas, A. A. and Onarinde, B. A. (2021) [46]	To measure the impact of COVID-19 on the eating and purchasing behaviours of people living in England based on sociodemographic variables	An online survey using questionnaires	911 (77 = Asian/Asian British, 38 = Black/Caribbean, 20 = mixed, 638 = White, 8 = Arab, 11 = others)	<ul style="list-style-type: none"> -BAME groups had the greatest effect of COVID-19 on decisionmaking and purchasing of healthier foods compared to participants from white backgrounds - Among the BAME groups, a considerable number of people from mixed or multiple ethnic groups had the lowest number of participants who had their decisions on healthier food affected "moderately or a little bit"

Table 1 (continued)

S.no	Authors	Aim	Method	Participants (N)	Main findings
4	Hassan, S. M. et al. (2021) [48]	To understand better, perceptions of risk and responses to COVID-19 of members of the Muslim community living in the Northwest of England, and to understand the facilitators and barriers to adherence to restrictions and guidance measures	An in-depth qualitative study using interviews and focus group discussions	47	<ul style="list-style-type: none"> -There were changes reported in the overall health behaviours of the participants -Positive hygiene practices and social distancing were reported among some participants -Participants also described additional precautions they were taking to reduce risk of transmission, including wearing face masks/covering (well before this became mandatory), wearing gloves, using hand sanitisers and disinfecting food packaging before putting it away

may drink less because of religious prohibitions. Likewise, in a multi-national survey which included BAME sample, Ammar *et al.* [53] found that a reduction in binge drinking was the major dietary habit change during the COVID-19 lockdown. Although Ammar *et al.* [53] suggests that younger people drank during the COVID-19 lockdowns due to reduced social interaction, Garnett *et al.* [45] reports that drinking more was independently associated with being younger. Furthermore, this scoping review reports that psychological stress projected changes in drinking behaviours. This stress could be associated with fear of contracting COVID-19 and becoming severely ill or low finances due to the associated economic loss. While stress is a risk factor for the inception of alcohol misuse, it can also act as a polarising factor for people to reduce alcohol intake and improve health [54].

Changes in eating and purchasing of healthier foods among BAME group was reported by one of the studies we reviewed [46]. Although the reason for the changes is not clear, it may be due to cultural, social or economic factors [55]. Pietrobelli *et al.* [56] identified an increase in the consumption of fruits among 41 children in Italy. This is similar in four other studies [50, 57–59]. This can be attributed to an increase in home cooking due to lockdown restrictions as well as WHO guidelines on the consumption of fruits and vegetables during lockdown [40, 60]. However, other studies globally have shown decrease in the consumption of fresh foods [40, 61]. A survey in Zimbabwe attributed this decrease to increased price and unavailability of fresh foods due to lockdown restrictions [49]. Another study in India reported that 32% of their respondents had indicated that an increase in price was a reason for their reduced intake of fruits and vegetables [62]. Nonetheless, there are very limited empirical evidence-based studies measuring the effects of COVID-19 on the dietary behaviours among BAME group in the UK.

This scoping review has further revealed the increase in protective health behaviours such as positive hygiene practices and social distancing. A possible explanation for these behaviours could be a means to mitigate risk of contracting COVID-19. Several studies have established links between risk-perceptions and protective health behaviours during pandemics. A study in Italy that explored the association between risk perceptions and compliance with recommendations during the 2009 Influenza H1N1 pandemic, reported that participants complied with recommended behaviours due to their perceived risk of contracting the virus [63]. A review of demographic and attitudinal determinants of protective behaviours during a pandemic further revealed that higher levels of perceived risk and severity of disease are

associated with adoption of recommended behaviours in a pandemic [64]. Conversely, an American study has found out that, while perceived risk of contracting COVID-19 may increase the level of protective health behaviours, perceived severity of COVID-19 did not [65]. Furthermore, an international study has found out that perceived risk of COVID-19, perceived severity of COVID-19 and trust in government were of little importance in voluntary compliance of protective health behaviours.

Nevertheless, this scoping review has shown that there is scarce evidence on health behaviour changes among BAME groups in terms of changes in health behaviours due to the COVID-19 pandemic. Smoking, high alcohol consumption, physical inactivity and a poor diet are four principal behavioural risks to health with the latter two also causing obesity [66]. Their prevalence varies across the population, although they tend to cluster in more deprived communities [33]. Also, evidence shows that these behavioural risks accounts for two-thirds of the incidence of cardiovascular diseases, chronic conditions, diabetes and cancer [67–69]. Further, more than 38 million deaths globally have been attributed to chronic conditions [70]. The existing inequalities on ethnic minorities following the COVID-19 pandemic, is now well reported. Available evidence has suggested that the cause of these inequalities is an intersection of deprivation, environmental, cultural, behavioural and physiological factors [33]. COVID-19 has highlighted the health inequalities experienced by ethnic minorities, thus, there is an urgent need to prevent and manage ill health in ethnic minority communities.

Strengths and limitations of the study

To the best of our knowledge, this is the first scoping review that aimed to explore the impact of COVID-19 on the changes in health behaviours among BAME groups in the UK. The search strategy of this present scoping review was constrained to articles published in English language due to the UK context. Also, this scoping review did not appraise the quality of the evidence in the primary studies like systematic reviews. As a result, the validity and methodological quality of the included studies are not known. Further, only four studies met the inclusion criteria and were included for this scoping review. Consequently, findings from this scoping review are very limited and may not be generalised. Therefore, there is need for further research to explore the experiences and health behaviour changes among ethnic minorities in the UK following the recent COVID-19 pandemic.

Conclusion

The COVID-19 pandemic had a significant impact on the health behaviour of BAME groups especially during the lockdowns. They reported effects such as changes in eating and purchasing of healthier foods, and low levels of physical activities. Hence, there is need to promote health awareness among BAME groups to encourage healthy living. In addition, programmes such as physical fitness activities that favour BAME groups can be put in place, for example BAME women's walking groups, men's walking groups and young people's exercise groups to encourage people from BAME backgrounds to engage in physical activities. Healthy food programmes, such as giving out healthy food vouchers and parcels can be distributed to people from BAME backgrounds who are not able to afford healthy food due to the impact of the COVID-19 pandemic. However, the COVID-19 pandemic increased positive hygiene among BAME groups which is important in preventing other diseases and infections. This scoping review has highlighted that there is limited evidence on the impact of COVID-19 on health behaviour focusing on BAME communities. Hence, there is a need for researchers to explore COVID-19 related health behaviour changes among BAME communities.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12889-023-15978-0>.

Additional file 1.

Acknowledgements

The authors are grateful for the support of the Institute of Health Research (IHR) at the University of Bedfordshire and the Luton Borough Council.

Authors' contributions

IN: Conceptualisation, Literature search, search term development, screening, analysis and writing (original draft, review and editing). NA: Conceptualisation, supervision, writing (review and editing). EC: Writing (review). IK: Writing (Review). DA: Literature search and search term development. GR: Writing (review). SC: Supervision. CE: Conceptualisation, review and supervision. PO: Writing (review). All authors read and approved the final manuscript.

Funding

National Institute for Health Research.

Availability of data and materials

All data generated or analysed during this study are included in this published article [and its supplementary information files].

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Institute for Health Research, University of Bedfordshire, Luton, UK. ²School of Psychology, University of Bedfordshire, Luton, UK. ³Luton Borough Council, Luton, UK.

Received: 8 August 2022 Accepted: 24 May 2023

Published online: 31 July 2023

References

- Naughton F, et al. Health behaviour change during the UK COVID-19 lockdown: findings from the first wave of the C-19 health behaviour and well-being daily tracker study. *Br J Health Psychol.* 2021;26:624–43.
- Short ES, Molburn S. Social determinants and health behaviors: conceptual frames and empirical advances. *Curr Opin Psychol.* 2015;5:78–84.
- McBride E, Chilcot J. The impact of COVID-19 on health behaviour, well-being, and long-term physical health. *Br J Health Psychol.* 2021;26:259–70.
- Badr H, et al. Psychosocial and health behavioural impacts of COVID-19 pandemic on adults in the USA: protocol for a longitudinal cohort study. *BMJ Open.* 2020;10(12):e044642.
- Rizzuto D, Fratiglioni L. Lifestyle factors related to mortality and survival: a mini review. *Gerontology.* 2014;60:327–35.
- WHO Global status report on noncommunicable diseases 2010, Geneva. 2011. Available at: <https://apps.who.int/iris/handle/10665/44579> (Accessed: 21 May 2022).
- Swann C, et al. Health systems and health-related behaviour change: a review of primary and secondary evidence' (Online). 2008 Available at: <https://www.nice.org.uk/media/default/About/what-we-do/NICE-guidance/NICE-guidelines/Public-health-guidelines/Additional-publications/Special-report-health-systems-and-health-related-behaviour-change.pdf>.
- Khaw K, et al. Combined Impact of health behaviours and mortality in men and women: the epic-norfolk prospective population study. *PLoS Med.* 2008;5(1): e12.
- GOV.UK. Living safely with respiratory infections, including COVID-19. 2021. Available at: <https://www.gov.uk/guidance/living-safely-with-respiratory-infections-including-covid-19>.
- Jones D, Neal RD, Duffy SRG, Scott SE, Whitaker KL, Brain K. Impact of the COVID-19 pandemic on the symptomatic diagnosis of cancer: the view from primary care. *Lancet Oncology.* 2020;21:748–50. [https://doi.org/10.1016/S1470-2045\(20\)30242-4](https://doi.org/10.1016/S1470-2045(20)30242-4).
- NHS England. Help us help you: NHS urges public to get care when they need it. 2020 Retrieved from <https://www.england.nhs.uk/2020/04/help-us-help-you-nhs-urges-public-to-get-care-when-they-need-it/>.
- Brown A, et al. Negative impact of the first COVID-19 lockdown upon health-related behaviours and psychological wellbeing in people living with severe and complex obesity in the UK. *Eclinical Medicine.* 2021;34: 100796.
- Rogers J. P. et al. 'Neurology and neuropsychiatry of COVID-19: a systematic review and meta-analysis of the early literature reveals frequent CNS manifestations and key emerging narratives', *J Neurol Neurosurg Psychiatry.* 2021;0, pp.1–10.
- Spence JC, Rhodes RE, McCurdy A, Mangan A, Hopkins D, Mummery WK. Determinants of physical activity among adults in the United Kingdom during the COVID-19 pandemic: the DUK-COVID study. *Br J Health Psychol.* 2021;26(2):588–605. <https://doi.org/10.1111/bjhp.12497>.
- Deschasaux-Tanguy M, et al. 'Diet and physical activity during the COVID-19 lockdown period (March-May 2020): results from the French NutriNet-Sante cohort study', *MedRxiv.* 2020 <https://doi.org/10.1101/2020.06.04.20121855>.
- McAtamney K, Mantzios M, Egan H, Wallis DJ. Emotional eating during COVID-19 in the United Kingdom: Exploring the roles of alexithymia and emotion dysregulation. *Appetite.* 2021;161:105120.
- Institute for Employment Studies. IES Working at Home Wellbeing Survey - Institute for Employment Studies (IES). 2020 Retrieved from <http://www.employment-studies.co.uk/resource/ies-working-home-wellbeing-survey>.

18. Rolland B, et al. 'Global changes and factors of increase in caloric/salty food intake, screen use, and substance use during the early covid-19 containment phase in the general population in France: a survey study. *JMIR Public Health Surveill.* 2020;6(3).
19. Evans J, Middlehurst C, Nilsson P. Smokers stock up on tobacco and nicotine products. 2020. Retrieved from: <https://www.ft.com/content/362d7d51-6561-493b-a29b-72784157cca7>. Accessed 01 June 2023.
20. Klemperer EM, et al. Change in tobacco and electronic cigarette use and motivation to quit in response to COVID-19. *Nicotine Tob Res.* 2020;22:1662–3. <https://doi.org/10.1093/ntr/ntaa072>.
21. Action on Smoking and Health. A million people have stopped smoking since the COVID pandemic hit Britain. 2020. Retrieved from <https://ash.org.uk/media-and-news/press-releases-mediaand-news/pandemicmillion/>.
22. Qin F, et al. Physical activity, screen time and emotional wellbeing during the 2019 novel Coronavirus outbreak in China. *Int J Environ Res Public Health.* 2020;17(14):5170.
23. Benatti FB, Ried-Larsen M. The effects of breaking up prolonged sitting time: a review of experimental studies. *Med Sci Sports Exerc.* 2015;47(10):2053–61.
24. Chandrasekaran B, Ganesan TB. Sedentarism and chronic disease risk in COVID-19 lockdown: a scoping review. *Scott Med J.* 2021;66(1):3–10.
25. Dempsey PC, Owen N, Yates TE, Kingwell BA, Dunstan DW. Sitting less and moving more: Improved glycaemic control for type 2 diabetes prevention and management. *Curr DiabRep.* 2016;16(11):114. <https://doi.org/10.1007/s11892-016-0797-4>.
26. Noble N, Paul C, Turon H, Oldmeadow C. Which modifiable health risk behaviours are related? a systematic review of the clustering of Smoking, Nutrition, Alcohol and Physical activity ('SNAP') health risk factors. *Prev Med.* 2015;81:16–41. <https://doi.org/10.1016/j.ypmed.2015.07.003>.
27. Abrams EM, Szefer SJ. COVID-19 and the impact of social determinants of health. *Lancet Respir Med.* 2020;8:659–61. [https://doi.org/10.1016/S2213-2600\(20\)30234-4](https://doi.org/10.1016/S2213-2600(20)30234-4).
28. The Health Foundation. Health equity in England: The marmot review 10 years on. 2020 Retrieved from <https://www.health.org.uk/publications/reports/the-marmot-review-10-yearson>.
29. Breakwell GM, et al. The COVID-19 Preventive behaviors index: development and validation in two samples from the United Kingdom. *Eval Health Prof.* 2021;44(1):77–86.
30. Robertson M, et al. Exploring changes in body image, eating and exercise during the COVID-19 lockdown: a UK survey. *Appetite.* 2020;159: 105062.
31. Aspinall PJ. 'Ethnic/Racial Terminology as a Form of Representation: A Critical Review of the Lexicon of Collective and Specific Terms in Use in Britain', *Genealogy*, 4(3), 87. Available at: 2020. <https://doi.org/10.3390/genealogy4030087>.
32. Ali N, Butt J. and Phillips M. Improving responses to the sexual abuse of Black, Asian and minority ethnic children. 2021 Available at: <https://www.csacentre.org.uk/documents/improving-responses-csa-of-black-asian-minority-ethnic-children/>.
33. Raleigh V. and Holmes J. The health of people from ethnic minority groups in England. 2021 Available at: <https://www.kingsfund.org.uk/publications/health-people-ethnic-minority-groups-england>.
34. Herle M, et al. Trajectories of eating behaviour during COVID-19 lockdown: longitudinal analyses of 22,374 adults. *Clin Nutr.* 2021. <https://doi.org/10.1016/j.clnesp.2021.01.046>.
35. Bann D. et al. 'Changes in the behavioural determinants of health during the COVID-19 pandemic: gender, socioeconomic and ethnic inequalities in five British cohort studies'. *J Epidemiol Community Health.* 2021;0:1–7.
36. Mostofsky E, et al. Key findings on alcohol consumption and a variety of health outcomes from the nurses' health study. *Am J Public Health.* 2016;106:1586–91. <https://doi.org/10.2105/AJPH.2016.303336>.
37. King's College London. Nearly a third of UK public drinking more alcohol than usual during the pandemic [Press release]. 2020 Retrieved from <https://www.kcl.ac.uk/news/nearly-a-third-of-uk-public-drinking-more-alcohol-than-usual-during-the-pandemic>.
38. Inman P. UK lockdown sends retail sales to record low, but alcohol sales rise. *The Guardian*: 2020 Retrieved from. <https://www.theguardian.com/business/2020/apr/24/uk-lockdown-sends-retail-sales-to-record-low-but-alcohol-sales-rise>.
39. Luthra R. Are there ethnic differences in adherence to recommended health behaviours related to Covid-19? 2021. Available at: <https://www.iser.essex.ac.uk/blog/2021/06/14/are-there-ethnic-differences-in-adherence-to-recommended-health-behaviours-related-to-covid-19>.
40. Bennett G, et al. The impact of lockdown during the COVID-19 outbreak on dietary habits in various population groups: a scoping review. *Front Nutr.* 2021;8: 626432.
41. Kumar K, et al. COVID-19 and ethnicity: challenges in rheumatology. *Rheumatology.* 2020;59(8):1802–3.
42. Zajacova A, et al. Changes in health behaviours during early COVID-19 and socio-demographic disparities: a cross-sectional analysis. *Can J Public Health.* 2020;111:953–62.
43. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol.* 2005;8(1):19–32.
44. Henderson LK, et al. How to write a cochrane systematic review. *Nephrology (Carlton).* 2010;15(6):617–24.
45. Garnett C, et al. Factors associated with drinking behaviour during COVID-19 social distancing and lockdown among adults in the UK. *Drug Alcohol Depend.* 2021;219: 108461.
46. Ogundijo DA, et al. Exploring the impact of COVID-19 pandemic on eating and purchasing behaviours of people living in England. *Nutrients.* 2021;13(5):1499.
47. Wickersham A, et al. Changes in physical activity among United Kingdom University students following the implementation of coronavirus lockdown measures. *Int J Environ Res Public Health.* 2021;18(6):2792.
48. Hassan SM, et al. How do Muslim community members perceive Covid-19 risk reduction recommendations - a UK qualitative study? *BMC Public Health.* 2021;21:449.
49. Matsungo TM, Chopera P. Effect of the COVID-19-induced lockdown on nutrition, health and lifestyle patterns among adults in Zimbabwe. *BMJ Nutrition, Prevention & Health.* 2020;3(2):205–12.
50. Rodriguez-Perez C, et al. Changes in dietary behaviours during the COVID-19 outbreak confinement in the Spanish COVIDiet study. *Nutrients.* 2020;12(6):1730.
51. Phillipou A, et al. Eating and exercise behaviors in eating disorders and the general population during the COVID-19 pandemic in Australia: initial results from the COLLATE project. *Int J Eat Disord.* 2020;1(10):1002.
52. Gallo LA, et al. The impact of isolation measures due to COVID-19 on energy intake and physical activity levels in Australian university students. *Nutrients.* 2020. <https://doi.org/10.1101/2020.05.10.20076414>.
53. Ammar A, et al. Effects of COVID-19 Home Confinement on Eating Behaviour and Physical Activity: Results of the ECLB-COVID19 International Online Survey. *Nutrients.* 2020;12(6):1583.
54. Jose BS, et al. Stressors and alcoholic consumption. *Alcohol Alcohol.* 2000;35(3):307–12.
55. Martin CA, et al. Socio-demographic heterogeneity in the prevalence of COVID-19 during lockdown is associated with ethnicity and household size: results from an observational cohort study. *EClinical Medicine.* 2020;25: 100466.
56. Pietrobelli A, et al. Effects of COVID-19 lockdown on lifestyle behaviors in children with obesity living in Verona, Italy: a longitudinal study. *Obesity (Silver Spring, Md).* 2020;28(8):1382–5.
57. Allabadi H, et al. Impact of COVID-19 lockdown on dietary and lifestyle behaviours among adolescents in Palestine. *Dynamics of Human Health.* 2020;7(2):2170.
58. Bhutani S, Cooper JA. COVID-19-related home confinement in adults: weight gain risks and opportunities. *Obesity (Silver Spring).* 2020;28(9):1576–7.
59. Ruiz-Roso MB, et al. COVID-19 lockdown and changes of the dietary pattern and physical activity habits in a cohort of patients with type 2 diabetes mellitus. *Nutrients.* 2020;4(12):2327.
60. WHO. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. 2020 Available at: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>.
61. Bracale R, Vaccaro CM. Changes in food choice following restrictive measures due to Covid-19. *Nutr Metab Cardiovasc Dis.* 2020;30:1423–6.
62. Mehta P, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *The Lancet.* 2020;395(10229):1033–4.
63. Prati G, et al. Compliance with recommendations for pandemic influenza H1N1 2009: The role of trust and personal beliefs. *Health Educ Res.* 2011;26(5):761–9.

64. Bish A, Michie S. Demographic and attitudinal determinants of protective behaviours during a pandemic: a review. *J Health Psychol.* 2010;15(4):797–824.
65. Wise T, et al. Changes in risk perception and self-reported protective behaviour during the first week of the COVID-19 pandemic in the United States. *Royal Society Open Science.* 2020;7: 200742.
66. Williams E, Buck D. and Babalola G. What are health inequalities? 2020. Available at: <https://www.kingsfund.org.uk/publications/what-are-health-inequalities>.
67. Beaglehole R, et al. Priority actions for the non-communicable disease crisis. *Health Policy.* 2011;377(9775):1438–47.
68. Kearns P. Organisational learning and development. 2014. Available at: <https://www.taylorfrancis.com/books/mono/10.4324/9781315777320/organizational-learning-development-paul-kearns> (Accessed: 21 May 2022).
69. World Health Organisation cardiovascular diseases (CVDs). 2020. Available at: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)#:~:text=The%20most%20important%20behavioural%20risk%20factors%20of%20heart,glucose%2C%20raised%20blood%20lipids%2C%20and%20overweight%20and%20obesity](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)#:~:text=The%20most%20important%20behavioural%20risk%20factors%20of%20heart,glucose%2C%20raised%20blood%20lipids%2C%20and%20overweight%20and%20obesity).
70. Ng K, et al. Physical activity and screen time among children and adolescents in Kazakhstan. *EQOL Journal.* 2019;11(2):19–27.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions



Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com