



**University of
Sunderland**

Onchonga, David, Omwoyo, Joshua and Nyamamba, Duke (2020) Assessing the prevalence of self-medication among healthcare workers before and during the 2019 SARS-CoV-2 (COVID-19) pandemic in Kenya. *Saudi Pharmaceutical Journal*, 28 (10). pp. 1149-1154. ISSN 1319-0164

Downloaded from: <http://sure.sunderland.ac.uk/id/eprint/16656/>

Usage guidelines

Please refer to the usage guidelines at <http://sure.sunderland.ac.uk/policies.html> or alternatively contact sure@sunderland.ac.uk.



Original article

Assessing the prevalence of self-medication among healthcare workers before and during the 2019 SARS-CoV-2 (COVID-19) pandemic in Kenya

David Onchonga^{a,b,*}, Joshua Omwoyo^b, Duke Nyamamba^b^a Doctoral School of Health Sciences, Faculty of Health Sciences, University of Pécs, Hungary^b Ministry of Health- Samburu County, Kenya

ARTICLE INFO

Article history:

Received 8 June 2020

Accepted 8 August 2020

Available online 14 August 2020

Keywords:

Self-medication

Prevalence

Healthcare workers

2019 SARS-CoV-2

Kenya

COVID-19

ABSTRACT

Background: Self-medication plays a key role in public health as it influences both negatively and positively on the health of individuals and the existing healthcare systems. This is especially the case during public health emergencies like the 2019 SARS-CoV-2 disease.

Objectives: The study aimed at assessing the prevalence of self-medication before and during the outbreak of COVID-19 pandemic among healthcare workers and its associated factors.

Methods: Stratified sampling was used to select 379 study respondents. The authors developed an online questionnaire, pretested and submitted to various online professional groups of different cadres of healthcare workers. Collected data was analysed using descriptive and inferential statistics. The data was presented in tables, graphs, percentages, and cross-tabulation with different variables. Multivariable logistic regression was used to ascertain factors that influence an individual's desire to self-medicate and associations were considered significant at $p < 0.05$.

Results: The overall prevalence of self-medication increased from 36.2% ($n = 137$) before the pandemic to 60.4% ($n = 229$) during the pandemic. The respondents' gender, level of education, age, marital status, participation in physical activity, and drug reaction events, were significantly associated with self-medication before and during the outbreak ($P < 0.05$). Multivariable logistic regression analyses showed that participants engaging in physical activities, working during the day and being healthy were less likely to self-medicate ($p < 0.01$).

Conclusion: Self-medication is an important health issue, especially during the COVID-19 pandemic. Continuous awareness creation and sensitization could help in reducing self-medication practices among healthcare workers.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Self-medication also known as self-care and/or self-administration of medication is a common practice across the world (Hughes et al., 2001). It involves situations where individuals, families, and/or communities take medicines to treat health conditions or symp-

toms without prescription or diagnosis from a qualified healthcare professional (Adolfo et al., 2000; Bennadi, 2014; Ruiz, 2010). Although self-medication has been practiced in many generations (Huffman, 2001), it is becoming a significant area of interest in healthcare research as it gives individuals with minor ailments freedom to make independent choices on how to manage their own health and wellbeing (Hughes et al., 2001). Research has demonstrated that there are a number of merits attributed to self-medication, such as improved access to medicines, especially over-the-counter drugs, which eases the burden levied to physicians and pharmacists in dealing with minor illnesses (Powis et al., 1996).

Studies have shown that although self-medication empowers individuals and communities to take control of their wellness, it has been associated with medical challenges (Mandal, 2015), which may have adverse impacts on their lives. This has been attributed to uninformed acts such as incorrect dosage, incorrect

* Corresponding author at: Doctoral School of Health Sciences, Faculty of Health Sciences, University of Pécs-Hungary, Pécs, Vörösmarty Mihály utca 4, 7621, Hungary.

E-mail addresses: onchonga.david@etk.pte.hu (D. Onchonga), joshu2064@gmail.com (J. Omwoyo), nyamamba1@gmail.com (D. Nyamamba).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

diagnosis of a particular health condition and improper route of administration of the self-medicated drugs. In addition, they may be using medicines for longer than the recommended time by the manufacturers, with a possibility of developing risk of dependence leading to abuse, improper storage of drugs, inappropriate choice of drug therapy, and increased prevalence of pathogenic resistance to drugs as some of the adverse effects (Menary et al., 2011).

Consumption of drugs without a prescription is triggered by factors such as the availability of drugs, easy access to drugs without time limits, improved supply of different varieties of medicines at affordable prices, and convenience in access to drugs compared to seeking treatment in the healthcare facilities (Smith, 1996).

A recently published google trends study on the interest in self-medication during COVID-19 pandemic indicated an upward trend. The study showed an increase in the number of searches for self-medication worldwide since the pandemic was declared, and this would be an indication of increased interests in self-medication across the globe (Onchonga, 2020).

Self-medication is commonly practised in countries where human resources for health are constrained. Most self-medication practices are triggered by relatively long waiting time in the health facilities, difficulties in securing an appointment with physicians, frequent closures of health facilities, recurrent stock-outs of essential medicines, and delays in getting treatment during emergencies (Parulekar et al., 2016).

For healthcare workers, research has revealed that the circumstances and the environment in which they work predisposes them to accessing drugs during their routine work. Although majority of the healthcare workers may be knowledgeable about the dangers posed as a result of self-medication, most of them find solace in self-medication especially when faced with work-related stress, pressure, discomfort, and anxiety (Omolase et al., 2011; Rotenberg, 2009).

Currently, the world is struggling with COVID-19 pandemic (Lai et al., 2020; Weiss and Leibowitz, 2011; Zu et al., 2020), which has been declared a global public health emergency (WHO, 2020). Due to its fast spread, the disease has threatened lives and caused psychological distress and anxiety to both healthcare workers and the general public in many parts of the world (Holmes, 2003). This situation has complicated the already fragile healthcare systems in resource-limited settings, especially those offering maternal and child health services (Onchonga et al., 2020a, 2020b). The focus has shifted to combating the pandemic; thereby curtailing the strides made in other healthcare indicators.

The current situation has been exacerbated by inadequacies in healthcare infrastructure, essential medical equipment in health facilities and inadequate health workforce. As a result, the available workers are forced to work for long hours, sometimes under difficult circumstances (Bogren et al., 2019), which prompts them to self-medicate to relieve the discomfort and anxiety attributed to burnouts.

This cross-sectional study was designed to investigate the pattern of self-medication among healthcare workers before and during the 2019 COVID-19 pandemic. The objectives of the study included: (a) to evaluate the prevalence of self-medication reported by healthcare workers before and during the outbreak, (b) to identify the common conditions for self-medication among healthcare workers before and during the outbreak, and (c) to investigate the sociodemographic characteristics and health patterns associated with self-medication among different cadres of healthcare workers before and during the outbreak. It was envisaged that the results would inform healthcare policy makers on measures to put in place to improve the well-being of healthcare workers as they strive to save lives.

2. Methods

2.1. Study design

This was part of a larger study that was conducted in Kenya on the effect of COVID-19 pandemic. The study population consisted of 385 healthcare workers that were directly involved in the management and control of COVID-19 disease pandemic. The study employed stratified sampling. Various cadres of healthcare professionals were used as strata. An online questionnaire was developed, pretested, revised, and sent to different healthcare professionals' online groups and platforms such as WhatsApp, Messenger, and Facebook. From each stratum, data was collected through random sampling of the received questionnaires until the required sample was achieved. The shared questionnaire was made anonymous to ensure data confidentiality and reliability.

2.2. Data collection

A questionnaire was developed by the investigators, which contained respondents' sociodemographic characteristics such as age, marital status, gender, level of education, religion, and cadre. Moreover, it contained questions on the prevalence of self-medication before and during the outbreak, conditions treated for self-medication before and during the outbreak, reasons for preferring self-medication before and during the outbreak, respondent's attitude regarding self-medication, and knowledge and perceptions regarding self-medication before and during the outbreak. Adverse drug reaction experiences were also investigated.

2.3. Data analysis and presentation

Collected data was analysed using the Statistical Package Programme for Social Science (SPSS) version 23.0.0. by IBM. Data was analysed and summarized using descriptive and inferential statistics, and later presented in tables, graphs, percentages, and cross-tabulation. Multivariable logistic regression analyses were used to ascertain factors that influence an individual's desire to self-medicate, and associations were considered significant at $p < 0.05$.

2.4. Ethical approvals

The approval to conduct this research was obtained from the Jaramogi Oginga Odinga Teaching and Referral Hospital Ethical Review Committee (IERC/JOOTRH/209/20). All the respondents were well-versed on the study intentions and were required to consent before admission to the study.

3. Results

3.1. Sociodemographic characteristics of the respondents

Out of a targeted sample of 385 respondents, approximately 98.4% ($n = 379$) questionnaires were correctly filled and returned. A total of 14 cadres participated in the study where; 30.6% ($n = 116$) were nurses. Regarding the gender of participants, 51.5% ($n = 195$) were female and 91.3% ($n = 346$) were Christians. About 65.4% ($n = 248$) were married and 59.6% ($n = 226$) had a university degree as shown in Table 1.

Table 1
Socio-demographic characteristics of respondents before and during COVID-19 Outbreak.

Socio-demographic variable		N	(%)
Gender	Female	195	51.5
	Male	184	48.5
Education	College diploma	153	40.4
	University Degree	226	59.6
Age	18–30	128	33.8
	31–40	159	42.0
	41–50	72	19.0
	51–60	20	5.3
Religion	Christian	346	91.3
	Muslim	17	4.5
	Hindu	4	1.1
	Traditional	2	0.5
	No religion	10	2.6
Marital status	Single	131	34.6
	Married	248	56.4
Geographical location	Rural	99	21.6
	Peri-urban	112	29.6
	Urban	168	44.3
Place of work	Public Health Facility	260	68.6
	Private practice	75	19.8
	NGO	36	9.5
	Pharmaceutical company	8	2.1
Cadre	Public Health Officers	86	22.7
	Nursing Officer	116	30.6
	Clinical Officers	13	3.4
	Pharmacy/Technologists	67	17.7
	Medical Officers	22	5.8
	Community Health Assistants	16	4.2
	Nutritionists	10	2.6
	Laboratory Scientists/ Technicians	14	3.7
	Physiotherapist	5	1.3
	Medical Consultants	8	2.1
	Medical Registrars	9	2.4
	Radiologists	6	1.6
	Health Records Information	2	0.5
	Dentist/ Dental Technologists	5	1.3

3.2. Prevalence and reported reasons for self-medication

In this study, 50.9% (n = 193) reported having felt unwell in the last three weeks preceding the interview. Approximately 60.4% (n = 229) reported having taken medicines without a doctor's prescription in the last 21 days preceding the study. This was a 24.3% increase since those who had self-medicated before the COVID-19 outbreak were 36.1% (n = 137). The respondents were also asked about the conditions that necessitated self-medication. Approximately 42% (n = 139) mentioned headache and migraine, 15% (n = 51) indicated joint and muscle pains, and 13% (n = 43) mentioned running nose (Fig. 1). There were no differences on the conditions that necessitated self-medication before and during the outbreak.

3.3. Perceptions of the respondents regarding their health status

About 19.8% (n = 75) of the respondents reported to be in perfect health, 74.7% (n = 238) had good health, and 5.5% (n = 21) had average or poor health status. Just about 90% (n = 341) of the respondents did not have any known psychiatric conditions, while 77.6% (n = 294) were actively participating in physical activities. Sleeping patterns were also investigated. About 14.8% (n = 56) of the respondents reported sleeping for more than eight hours a day, 79.9% (n = 303) had a sleeping pattern of

between five and eight hours a day, and 5.3% (n = 20) had a sleeping pattern of below four hours a day. There were no differences in sleeping patterns before and during the outbreak.

3.4. Relationship between sociodemographic characteristics and self-medication

Chi-square test was conducted to compare self-medication before and during the outbreak with sociodemographic variables. The results showed statistically significant differences between self-medication and gender, physical activity, adverse drug reaction, work shift, marital status, level of education, and cadre.

There was no statistical significance between self-medication and residency of respondents, health status, psychiatric condition, place of work, and sleeping patterns (Table 2).

3.5. Regression analysis of the factors prompting respondents to self-medication

The study applied multivariable logistic regression analyses to ascertain factors influencing an individual's desire to self-medicate. The respondents that undertook physical activity, worked on day shift and were healthy; were less likely to self-medicate ($p < 0.01$). Variables such as gender, drug reaction, reading drug instructions before use, level of education, having a known psychiatric condition, and perceived health status did not correlate with self-medication (Table 3).

3.6. Respondent's knowledge and practices regarding self-medication before and during the outbreak

The number of respondents who read the manufacturer's manual/leaflet all the time before purchasing medicines increased from 58.8% (n = 223) before the outbreak to 72% (n = 273) while those who did not read at all reduced from 5.1% (n = 19) before the outbreak to 1% (n = 4) during the outbreak. About 70.4% (n = 267) of the respondents noted that self-medication was not safe. Regarding the route of administration of the purchased medicine, dosage, and side effects, there was an increase in knowledge from 69.4% (n = 263) before the outbreak to 75% (n = 284) during the outbreak, an indication of an increase in knowledge regarding the route of administration of the purchased medicines.

3.7. Consideration for self-medication

About 31.1% (n = 118) considered the brand of the medicine, 18.5% (n = 70), the pharmaceutical company manufacturing the medicine, while 17.4% (n = 66) considered the cost. Access to self-medication changed significantly before and during the outbreak. About 16.9% (n = 64) obtained the medicines from the primary health facility, 56.5% (n = 214) from pharmacy stores while 26.6% (n = 101) from hospital pharmacies before the outbreak, while, 89.5% (n = 340) obtained medication from pharmacy stores and only 10.5% (n = 39) from primary health facilities. None of the respondents obtained medication from hospital pharmacies during the outbreak.

3.8. Adverse drug reaction (ADR) following self-medication

Among the respondents who self-medicated, 22.4% (n = 85) had experienced perceived adverse drug reaction during the outbreak. There was a noteworthy association between self-medication and adverse drug reaction during the outbreak ($X^2 = 4.479$, $p = 0.034$). Further, there was a significant association between taking drugs for chronic illness and experiencing adverse drug reactions during the outbreak ($X^2 = 4.897$, $p = 0.02$), with results




Fig. 1. Conditions for self-medication before and during COVID-19 Outbreak.

Table 2
Relationship between Demographic Variables and Self-Medication before and during COVID-19 Outbreak.

Socio-demographic		Self-medication during COVID-19	Self-medication before COVID-19	χ^2	p value
		n (%)	n (%)		
1. Gender	Female	129(56.3)	66(44.0)	5.518	0.019
	Male	100(43.7)	84(56.0)		
2. Education	College Diploma	83(36.2)	70(46.7)	4.090	0.043
	University	146(63.8)	80(53.3)		
3. Age	18–30	69(30.1)	59(39.3)	10.592	0.014
	31–40	111(48.5)	48(32.0)		
	41–50	37(16.2)	35(23.3)		
	51–60	12(5.2)	8(5.4)		
4. Marital status	Single	119(52.0)	32(21.3)	36.606	0.000
	Married	110(48.0)	118(78.7)		
5. Geographical location	Rural	57(24.9)	42(28.0)	1.353	0.509
	Peri-urban	65(28.4)	47(31.3)		
	Urban	107(46.7)	61(40.7)		
6. Physical activity	Yes	181(79.0)	48(21.0)	9.578	0.002
	No	97(64.7)	53(35.3)		
7. Drug reaction event	Yes	135(59.0)	94(41.0)	14.759	0.000
	No	117(78.0)	33(22.0)		
8. Health status	Perfect health	38(16.6)	37(24.7)	3.860	0.145
	Good health	177(77.3)	106(70.7)		
	Average health	14(6.1)	7(4.7)		
9. Psychiatric condition	Yes	205(89.5)	24(10.5)	0.132	0.716
	No	136(90.7)	14(9.3)		
10. Sleeping pattern	More than 8 hrs	28(12.2)	28(18.7)	3.523	0.172
	5–8 hrs	187(81.7)	116(77.3)		
	1–4 hrs	14(6.1)	6(4.0)		

suggesting that those taking drugs for chronic illness during the outbreak were more likely to develop adverse drug reactions. Adverse drug effects prompted approximately 60.8% (n = 62) to stop self-medication (Table 4). The study did not find any significant association between self-medication and adverse drug reaction before the outbreak of COVID-19.

3.9. The opinion of respondents regarding self-medication and COVID-19 disease

Of the 379 respondents, 214 (64.3%) believed that the COVID-19 pandemic had necessitated self-medication. About 55.9% (n = 212)

reported that they would continue with self-medication post-COVID-19 pandemic. When asked about their opinion regarding self-medication and the outbreak, approximately 84.1% (n = 280) reported that the pandemic had increased the desire for self-medication for the general public. On a positive note, 91.3% (n = 304) would not advise other people to self-medicate.

4. Discussion

This study was conducted during the COVID-19 pandemic to investigate the prevalence of self-medication and associated factors among healthcare workers in Kenya before and during out-

Table 3

Multivariable logistic regression analyses on influencing factors of self-medication during the COVID-19 outbreak.

Independent variables (n = 379)	B	S.E.	Wald X2	P	Exp(B)	(95% CI)
Physical activity (Yes/No)	-1.160	0.356	10.615	0.001	0.313	(0.156,0.630)
Gender (Female/Male)	-0.081	0.308	0.069	0.793	0.922	(0.504,1.687)
Drug reaction event (Yes/No)	-0.469	0.338	1.929	0.165	0.625	(0.322,1.213)
Work shift (Night/Day)	-4.414	1.155	14.597	0.000	0.012	(0.001,0.117)
Reading prescription (All the time/Sometimes/Not at all)	-0.105	0.626	0.028	0.866	0.900	(0.264,3.070)
Level of education (Diploma/Degree)	0.890	0.340	6.832	0.009	2.434	(1.249,4.743)
Any known psychiatric condition (Yes/No)	0.169	0.519	0.106	0.745	1.184	(0.428,3.274)
Health status (Perfect /Good/Average)	0.701	0.716	0.960	0.327	2.016	(0.496,8.201)
Feeling unwell in the last 3 weeks (Yes/No)	-2.587	0.346	55.910	0.000	0.075	(0.038,1.48)
Constant	-20.099	40193.683	0.000	1.000	0.000	

Table 4

Action taken following adverse drug reaction event during COVID-19 outbreak.

Action taken after adverse drug reaction event	N	(%)
Went to a private doctor	12	11.8
Went to primary health centre	17	16.7
Went to a pharmacist	11	10.8
Stopped taking the medicine	62	60.8

break. The prevalence of self-medication had increased from 36.2% before the outbreak to 60.4%, during the outbreak. These results were close to the findings from a similar study undertaken in Nigeria (Bamgboye et al., 2006). In the current study, an analysis was done to determine the prevalence of self-medication across different cadres of healthcare workers, and the nursing officers reported a prevalence of 32.8% during the outbreak. This finding is similar to a study conducted in Brazil on the prevalence of self-medication among nurses (Rotenberg, 2009).

In the current study, several considerations for the preference of particular drugs for self-medication were evaluated. The brand of the medicine, cost, and the manufacturing company were the primary considerations. This, too, has been found in similar studies (Selvaraj and Ramalingam, 2014). The majority of the respondents took medicines for pain relief, more specifically headaches, migraine, fever, joint and muscle pains before and during the outbreak. Similar conditions have been documented as a common reason for self-medication (Auta et al., 2012).

Self-medication was significantly associated with the level of education as those with degrees were more likely to self-medicate both before and during the outbreak. These findings agree with similar studies (López and Dennis, 2009), and this has been attributed to the vast knowledge on the medication as most medical graduates do possess a higher level of understanding of the over-the-counter drugs, including their prescription and side effects. Of note was that some studies have given divergent results regarding this finding. A similar study (Sharma et al., 2005) reported that self-medication was higher among respondents who had secondary education and below compared to those with advanced education.

In the current study, gender was statistically significant. More female participants were likely to self-medicate before and during the outbreak. However, similar studies from other regions (Bennadi, 2014; Shankar et al., 2002) have shown that male healthcare workers are more likely to self-medicate than their female counterparts (Belachew Gutema et al., 2011; Sawalha, 2007). In this study, participants who engaged in physical activity were less likely to self-medicate both before and during the outbreak. This finding agrees with existing evidence on physical activity and self-medication (Wijesinghe et al., 2012).

The study participants were all working in healthcare institutions where access to medical care was guaranteed as all the respondents had a medical insurance cover. However, this did

not deter them from preferring self-medication. This study agrees with similar studies that have shown that open access to healthcare does not significantly reduce self-medication (Osaboro et al., 2013).

In addition, it was noted that work shifts triggered self-medication; those working at night were more likely to self-medicate both before and during the outbreak. Similar studies have indicated that most healthcare workers on night shift tend to self-medicate as a result of unfavourable working conditions, and working for long hours with minimum rest, which may contribute to lassitude (Rotenberg, 2009).

The findings from multivariable logistic regression analyses showed that the respondents who undertook physical activity, worked during the day, and those who were healthy, correlated well with the prevalence of self-medication ($p < 0.01$). Similar studies conducted in other regions have shown similar results (Lei et al., 2018).

5. Study limitations

This study was restricted to healthcare workers who were currently actively involved in managing the COVID-19 pandemic and had self-medicated 21 days preceding the interview. Therefore, there is a possibility of recall bias. Also, cross-sectional studies do not permit an establishment of a clear cause and effect.

6. Conclusion

Self-medication is a significant health issue, especially during COVID-19 pandemic, and a refresher course in the form of continued medical education (CME) should be encouraged. Awareness creation is also needed, not only to the healthcare workers but also to the general public; especially on the risks involved since the potential danger of self-medication might be underestimated during emergencies. There is need to undertake a study on how ethnic differences, and religious beliefs would influence self-medication among healthcare workers.

Funding

This study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors

CRedit authorship contribution statement

David Onchonga: Conceptualization, Methodology, Investigation, Data curation, Writing - original draft, Supervision. **Joshua Omwoyo:** Visualization, Investigation, Software, Validation, Writing - original draft. **Duke Nyamamba:** Writing - review & editing, Formal analysis, Writing - original draft.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Adolfo, F., Caamano, F., Gestal-Otero, J.J., 2000. Sociodemographic factors related to self-medication in Spain. *Eur. J. Epidemiol.* 16, 19–26. <https://doi.org/10.1023/A:1007608702063>.
- Auta, A., Omale, S., Folorunsho, T.J., David, S., Banwat, S.B., 2012. Medicine vendors: Self-medication practices and medicine knowledge. *N. Am. J. Med. Sci.* 4 (1), 24–28. <https://doi.org/10.4103/1947-2714.92899>.
- Bamgboye, E.A., Amoran, O.E., Yusuf, O.B., 2006. Self medication practices among workers in a tertiary hospital in Nigeria. *Afr. J. Med. Med. Sci.* 35 (4), 411–415. 17722805.
- Belachew Gutema, G., Alemayehu Gadisa, D., Fikadu Berhe, D., Hadgu Berhe, A., Ghezu Hadera, M., Solomon Hailu, G., Gebresamuel Abrha, N., Yarlagaadda, R., Wondimu Dagne, A., Abebe Kidanemariam, Z., 2011. Self-medication practices among health sciences students: The case of Mekelle University. *J. Appl. Pharm. Sci.* 1 (10), 183–189.
- Bennadi, D., 2014. Self-medication: A current challenge. *J. Basic Clin. Pharm.* 5 (1), 19–23. <https://doi.org/10.4103/0976-0105.128253>. 24808684.
- Bogren, M.U., Erlandsson, K., Byrskog, U., Bogren, M., Johansson, A., Kalid, M., Igal, A. A., Mohamed, J., Said, F., Pedersen, C., Osman, F., 2019. Health workforce perspectives of barriers inhibiting the provision of quality care in Nepal and Somalia-A qualitative study. *Sex. Reprod. Healthc.* 23, <https://doi.org/10.1016/j.srhc.2019.100481> 100481.
- Holmes, K.V., 2003. SARS coronavirus: a new challenge for prevention and therapy. *J Clin Invest.* 111 (11), 1605–1609. <https://doi.org/10.1172/JCI18819>.
- Osaboro, E., Clement, C.A., Adebola, O.E., Ejike, B.E., Laura, E., Ikechukwu, U.M., 2013. Self-medication practices among dental, midwifery and nursing students. *Eur. J. Gen. Dent.* 2 (1), 54–57. <https://doi.org/10.4103/2278-9626.106813>.
- Huffman, M.A., 2001. Self-Medicative Behavior in the African Great Apes: An Evolutionary Perspective into the Origins of Human Traditional Medicine: In addition to giving us a deeper understanding of our closest living relatives, the study of great ape self-medication provides a window into the origins of herbal medicine use by humans and promises to provide new insights into ways of treating parasite infections and other serious diseases. *BioScience* 51 (8), 651–661. [https://doi.org/10.1641/0006-3568\(2001\)051\[0651:SMBITA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0651:SMBITA]2.0.CO;2).
- Hughes, C.M., McElnay, J.C., Fleming, G.F., 2001. Benefits and risks of self medication. *Drug Safety* 24, 1027–1037. <https://doi.org/10.2165/00002018-200124140-00002>.
- Lai, C.-C., Shih, T.-P., Ko, W.-C., Tang, H.-J., Hsueh, P.-R., 2020. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *Int. J. Antimicrob. Agents* 55, (3). <https://doi.org/10.1016/j.ijantimicag.2020.105924> 105924.
- Lei, X., Jiang, H., Liu, C., Ferrier, A., Mugavin, J., 2018. Self-medication practice and associated factors among residents in Wuhan, China. *Int. J. Environ. Res. Public Health* 15 (1). <https://doi.org/10.3390/ijerph15010068>.
- López, J.J., Dennis, R.M.S., 2009. A study of self-medication in a neighborhood in Bogotá. *Rev. Salud Pública* 11 (3) 432–442. URL: http://www.scielo.org.co/scielo.php?pid=S0124-00642009000300012&script=sci_arttext&lng=en (accessed 5.20.20)
- Mandal, S., 2015. Can over-the-counter antibiotics coerce people for self-medication with antibiotics? *Asian Pacific J. Trop. Dis.* 5 (1), S184–S186. [https://doi.org/10.1016/S2222-1808\(15\)60886-6](https://doi.org/10.1016/S2222-1808(15)60886-6).
- Menary, K.R., Kushner, M.G., Maurer, E., Thuras, P., 2011. The prevalence and clinical implications of self-medication among individuals with anxiety disorders. *J. Anxiety Disord.* 25 (3), 335–339. <https://doi.org/10.1016/j.janxdis.2010.10.006>.
- Omolase, C.O., Adeleke, O.E., Afolabi, A.O., Ofolabi, O.T., 2011. Self medication amongst general outpatients in a Nigerian community hospital. *Ann. Ibadan Postgrad. Med.* 5 (2), 64–67. <https://doi.org/10.4314/aipm.v5i2.64032>.
- Onchonga, D., 2020. A Google Trends study on the interest in self-medication during the 2019 novel coronavirus (COVID-19) disease pandemic. *Saudi Pharm. J.* 28 (7), 903–904. <https://doi.org/10.1016/j.jpsps.2020.06.007>. 32641903.
- Onchonga, D., MoghaddamHosseini, V., Keraka, M., Várnagy, Á., 2020. Prevalence of fear of childbirth in a sample of gravida women in Kenya. *Sex. Reprod. Healthc.* 24, 100510. <https://doi.org/10.1016/j.srhc.2020.100510>.
- Onchonga, D., Várnagy, Á., Keraka, M., Wainaina, P., 2020. Midwife-led integrated pre-birth training and its impact on the fear of childbirth. A qualitative interview study. *Sex. Reprod. Healthc.* 25, 100512. <https://doi.org/10.1016/j.srhc.2020.100512>.
- Parulekar, M., Mekoth, N., Ramesh, C.M., Parulekar, A., 2016. Self-medication in developing countries a systematic review. *J. Pharm. Technol. Res. Manage.* 4 (2), 103–127. <https://doi.org/10.15415/jptrm.2016.42007>.
- Rotenberg, A.R.R.B.R.H.G.L., 2009. Self-medication among nursing workers from public hospitals [WWW Document]. <https://doi.org/https://doi.org/10.1590/S0104-11692009000600014>
- Powis, M.G., Rogers, P.J., Wood, S.M., 1996. United Kingdom Community Pharmacists' Views on Recent "POM-to-P" Switched Medicines. *Soc. Admin. Pharm.* 13, 188–197.
- Ruiz, M., 2010. Risks of self-medication practices. *Curr. Drug Saf.* 5, 315–323. <https://doi.org/10.2174/157488610792245966>.
- Sawalha, A.F., 2007. Assessment of Self-Medication Practice among University Students in Palestine: Therapeutic and Toxicity Implications. *The Islamic University Journal* 15 (2).
- Selvaraj, K., Ramalingam, G.K.S.A., 2014. Prevalence of self-medication practices and its associated factors in Urban Puducherry. *India* 5 (1), 32–36 <https://doi.org/10.4103/2229-3485.124569>.
- Shankar, P.R., Partha, P., Shenoy, N., 2002. Self-medication and non-doctor prescription practices in Pokhara valley, Western Nepal: A questionnaire-based study. *BMC Fam. Pract.* 3 (17), 1–7. <https://doi.org/10.1186/1471-2296-3-17>.
- Sharma, R., Verma, U., Sharma, C.L., Kapoor, B., 2005. Self-medication among urban population of Jammu city. *Indian J. Pharmacol.* 37 (1), 40–43. <https://doi.org/10.4103/0253-7613.13856>.
- Smith, S.K., 1996. Over the counter drugs. *BMJ* 312, 593. <https://doi.org/10.1136/bmj.312.7031.593>.
- Weiss, S.R., Leibowitz, J.L., 2011. Coronavirus pathogenesis. *Advances in Virus Research*, 81. Academic Press Inc., pp. 85–164 <https://doi.org/10.1016/B978-0-12-385885-6.00009-2>.
- WHO, 2020. Coronavirus disease 2019 (COVID-19) Situation Report – 81.
- Wijesinghe, P.R., Jayakody, R.L., Seneviratne, R.D.A., 2012. Prevalence and predictors of self-medication in a selected urban and rural district of Sri Lanka. *J. Public Health* 1 (1), 28–41. doi: <https://apps.who.int/iris/handle/10665/329805>.
- Zu, Z.Y., Jiang, M.D., Xu, P.P., Chen, W., Ni, Q.Q., Lu, G.M., Zhang, L.J., 2020. Coronavirus Disease 2019 (COVID-19): A Perspective from China. *Radiology* 296 (2). <https://doi.org/10.1148/radiol.202000490>.