



**University of
Sunderland**

Walter, Fiona M, Emery, Jon D, Mendonca, Silvia, Hall, Nicola, Morris, Helen C, Mills, Katie, Dobson, Christina, Bankhead, Clare, Johnson, Margaret, Abel, Gary A, Rutter, Matthew D, Hamilton, William and Rubin, Greg P (2016) Symptoms and patient factors associated with longer time to diagnosis for colorectal cancer: results from a prospective cohort study. *British Journal of Cancer*, 115 (5). pp. 533-541. ISSN 0007-0920

Downloaded from: <http://sure.sunderland.ac.uk/id/eprint/6612/>

Please refer to the usage guidelines at <http://sure.sunderland.ac.uk/policies.html> or alternatively contact sure@sunderland.ac.uk.

Symptoms and patient factors associated with longer time to diagnosis for colorectal cancer: results from a prospective cohort study

Running title: Factors affecting diagnostic intervals for colorectal cancer

Authors' names, affiliations, full address, postcode:

Fiona M Walter (corresponding author- fmw22@medschl.cam.ac.uk)

Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Jon D Emery

Department of General Practice
University of Melbourne
Australia

Silvia Mendonca

Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Nicola Hall

At the time of completing this research:
School of Medicine, Pharmacy & Health
Durham University
Stockton on Tees TS17 6BH

Helen C Morris

At the time of completing this research:
Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Katie Mills

Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Christina Dobson

School of Medicine, Pharmacy & Health
Durham University
Stockton on Tees TS17 6BH

Clare Bankhead

Nuffield Department of Primary Care Health Sciences
University of Oxford
Oxford OX2 6GG

Margaret Johnson

Lay member, c/o:
Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Gary A Abel

Primary Care Unit
Department of Public Health & Primary Care
University of Cambridge
Cambridge CB1 8RN

Matthew D Rutter

School of Medicine, Pharmacy & Health
Durham University
Stockton on Tees TS17 6BH
University Hospital of North Tees
Stockton on Tees TS19 8PE

William Hamilton

College House, St Luke's Campus
University of Exeter
Exeter EX2 4TE

Greg P Rubin

School of Medicine, Pharmacy & Health
Durham University
Stockton on Tees TS17 6BH

Word count: Abstract 210 words, Main text 4000 words, Number of Tables: 6 (& supplemental table and figure)

Keywords: Symptoms, diagnosis, diagnostic interval, time to presentation, delays, helpseeking, stage, colorectal cancer, colon cancer, rectal cancer.

ABSTRACT

Background

To investigate symptoms, clinical factors and socio-demographic factors associated with colorectal cancer (CRC) diagnosis and time to diagnosis.

Methods

Prospective cohort study of participants referred for suspicion of CRC in two English regions. Data were collected using a patient questionnaire, primary care and hospital records. Descriptive and regression analyses examined associations between symptoms and patient factors with: Total Diagnostic Interval (TDI), Patient Interval (PI), Health System Interval (HSI), stage.

Results

2,677 (22%) participants responded; after exclusions, 2,507 remained. 6.1% were diagnosed with CRC (56% late stage), 2.0% with other cancers, 91.9% with no cancer. Half the cohort had a solitary first symptom (1332, 53.1%); multiple first symptoms were common. In this referred population rectal bleeding was the only initial symptom more frequent among cancer than non-cancer cases (34.2% vs 23.9%, $p=0.004$). There was no evidence of differences in TDI, PI or HSI for those with cancer versus non-cancer diagnoses (median TDI CRC 124 vs non-cancer 138 days, $p=0.142$). First symptoms associated with shorter TDIs were rectal bleeding, change in bowel habit, 'feeling different' and fatigue/tiredness. Anxiety, depression and gastro-intestinal comorbidities were associated with longer HSIs and TDIs. Symptom duration-dependent effects were found for rectal bleeding and change in bowel habit.

Conclusion

Doctors, as well as patients, respond less promptly to some symptoms of colorectal cancer than others. Healthcare professionals should be vigilant to the possibility of CRC in patients with relevant symptoms and mental health or gastro-intestinal comorbidities.

INTRODUCTION

Colorectal cancer (CRC) is the third most common cancer world-wide and the UK's fourth commonest (CancerResearchUK). Despite the introduction of screening programmes in many countries, most cases present symptomatically. Although five-year survival rates for CRC have improved to over 60% in the last decade, CRC patients in the UK continue to have poorer survival rates than those in comparable countries (Allemani et al., 2015, Coleman et al., 2011). This may partly be due to longer intervals between the onset of cancer symptoms and the patient's presentation to healthcare, compounded by unequal access to optimal diagnostics and specialist care, leading to more late stage diagnoses (Maringe et al., 2013, Neal and Allgar, 2005). There are also gender, age and region differences in time to diagnosis across the UK (Din et al., 2015).

The diagnostic process comprises a series of stages or intervals, each contributing to the overall period of time between onset of symptom/s and initiation of treatment (Weller et al., 2012). The Total Diagnostic Interval (TDI) comprises the Patient Interval (PI- from first symptom onset to first healthcare consultation) and the Health System Interval (HSI- from first consultation via referral and investigations to diagnosis and initiation of treatment) (Weller et al., 2012): shorter PIs and HSIs are associated with improved CRC prognoses (Neal et al., 2015, Topping et al., 2011). For CRC, recent evidence suggests that both intervals are longer than for other common cancers such as lung or urological cancer (Keeble et al., 2014, Neal et al., 2014a, Dregan et al., 2013, Lyratzopoulos et al., 2015b).

CRC patients may be symptomatic for many months before presentation. They may experience multiple symptoms, both bowel-specific (rectal bleeding, change in bowel habit, abdominal pain) and systemic (loss of weight or appetite, fatigue) (Rasmussen et al., 2015, Hamilton et al., 2005). Not all individuals interpret their initial symptoms as serious, and may attribute them to normal variation, aging, lifestyle or other comorbidities (N Hall, 2015, McLachlan et al., 2015, Macleod et al., 2009). International comparisons suggest that lower age-related risk and highest perceived barriers to symptomatic presentation were reported in the UK (Forbes et al., 2013); other potential influences include negative beliefs about cancer outcomes (Beeken et al., 2011) (Lyratzopoulos et al., 2015a) and poor awareness of the risk of cancer (Quaife et al., 2014, Simon et al., 2010).

Once the decision to seek medical help has been made, further delays may occur. One third of CRC patients have three or more consultations with a general practitioner (GP) prior to referral, compared with only 17.9% for all cancers (Lyratzopoulos et al., 2013). Furthermore, the pathway to diagnosis in primary care may be complex, with delays arising when presentation is complicated by comorbid conditions, when referral is delayed or declined, or from false-negative investigation (Rubin et al., 2014).

Symptoms of possible CRC much more commonly arise from benign or self-limiting conditions, and GPs face a considerable challenge to evaluate patients whilst making efficient use of hospital-based resources (Hansen et

al., 2015). Much of the evidence about the predictive value of symptoms for cancer or their association with later diagnosis comes from retrospective studies of people with CRC (Esteva et al., 2013) or from general practice datasets (Hamilton et al., 2005); these are limited by quality of data recording. Little is known about the diagnostic pathways of those whose symptoms transpire not to be caused by CRC or about which symptoms are associated with less timely diagnosis. We therefore recruited a prospective cohort of patients with symptoms suggestive of CRC at the point of their referral. We aimed to investigate the symptoms, clinical factors and socio-demographic factors associated with CRC diagnosis and time to diagnosis.

METHODS

Setting & governance

We recruited patients from four secondary care hospitals in the East (n=1) and North East of England (n=3) between December 2010 and March 2013. We gained all appropriate NHS ethics (reference: 10/H0306/50) and clinical governance approvals. The SYMPTOM colorectal study was conducted alongside the SYMPTOM lung and pancreas studies, collectively part of the NIHR-funded DISCOVERY programme of applied research. The methods of data collection and analysis have already been described with the reporting of the SYMPTOM lung study (Walter et al., 2015) so will be outlined briefly here.

Patient recruitment

All GP referral letters to urgent and routine colorectal and gastrointestinal clinics in participating hospitals were reviewed by a research nurse, in order to identify patients aged ≥ 40 years whose referral mentioned any one or more symptoms from a pre-specified list of symptoms known to be associated with CRC (Appendix A). These patients were sent an information sheet, the SYMPTOM colorectal questionnaire, and a reply-paid envelope for return. Patients were not eligible for the study if they were already undergoing treatment for any cancer (excluding non-melanotic skin cancer), or had such serious mental and/or physical disease that they were not considered suitable to complete a questionnaire, based on review of the medical record by the research nurse. Patients referred as a result of participation in the Bowel Cancer Screening Programme were excluded. No reminder letters were sent to non-responders, as required by the research ethics committee.

Data collection

We based our data collection, analysis and reporting on the Aarhus statement for the conduct of cancer diagnostic studies, and STROBE guidelines for reporting observational studies (von Elm et al., 2007, Weller et al., 2012).

Patient data

Patient data was collected using the SYMPTOM questionnaire, completed and returned to the research team within three months of being sent it; most completed it within 2 weeks and before receiving their diagnosis. The SYMPTOM colorectal questionnaire was derived from the previously-validated C-SIM questionnaire (Neal et al., 2014b). This had been developed for use among people recently diagnosed with cancer, and so we modified it for use before diagnosis, consulting widely with clinical and research colleagues and patient representatives to ensure it was worded sensitively, since cancer may not have been explicitly mentioned as a possibility by the referring GP. The questionnaire asked for the first symptom noticed by the participant, and then sought the presence or absence of specific symptoms (Appendix B). The remaining sections contained items about other symptoms and patient factors, including demographic characteristics and comorbidities.

Primary care data

Participants' GPs completed a short proforma from the patient's clinical record, including dates of the first presentation with any symptom from the questionnaire within the previous two years, plus its duration, if recorded.

Hospital data

We extracted data from hospital medical records on: date and type of referral (urgent, routine, emergency, other); date of first specialist consultation; dates of investigations and their findings; date of diagnosis (histological, clinical), and dates of MDT meetings and their clinical decisions. Data extraction took place a minimum of 6 months after recruitment to allow sufficient time for completion of investigation and initiation of treatment. We undertook double data abstraction of a 5% sample of selected hospital data (dates of referral, first appointment, diagnosis, stage) and confirmed an acceptable level of agreement (>80% for dates, >90% for diagnosis and stage, Cohen's Kappa>0.75 for all), with any discrepancies generally minor.

Data handling

Clinical outcomes

We used the date on the first confirmatory histology report as the date of diagnosis when this was available for both cancer (ICD codes C18-C20), and non-malignant conditions (Weller et al., 2012). Otherwise we used the first date of the clinical diagnosis in the hospital medical record. Participants were classified into three groups: those with primary CRC (CRC), other intra-abdominal cancer (OC) and no cancer (NC). The main analyses focused on CRC vs NC, with secondary analyses including all cancers (CRC plus OC) vs NC. CRC staging was categorised using TNM status at diagnosis (Travis et al., 2011), and further categorised into Early Stage (stages I and II) and Late Stage (stages III and IV). Complex diagnoses, or cases with incomplete data, were agreed by an adjudication group of clinicians associated with the study (FW, JE, GR, MR).

Demographic and clinical variables

Demographic details included: gender; age (treated as a continuous variable); ethnicity (coded as white vs non-white); smoking status; educational status; occupational status; living alone; and postcode, used to assign groups to quintiles of the Index of Multiple Deprivation (IMD) (1 'least deprived' to 5 'most deprived'). Clinical variables relating to comorbidities included: gastro-intestinal disease (inflammatory bowel disease, irritable bowel syndrome, peptic ulcer); respiratory disease (Chronic Obstructive Pulmonary Disease /asthma/other lung disease), anxiety/depression, heart disease, diabetes, arthritis. Family history of cancer was also identified (present vs absent).

Symptoms

All symptoms reported up to two years before diagnosis were entered into the analysis (symptoms of longer than 2 years duration were unlikely to be associated with the subsequent diagnosis) (Biswas et al., 2015) (Hamilton et al., 2005). Where participants provided estimated dates, these were converted to exact dates by adapting an algorithm used in the C-SIM trial (Neal et al., 2014b, Walter et al., 2015). We used exact dates where available and estimated dates otherwise. Furthermore, if a participant's unprompted, or free text, first symptom matched their response to a question about specific symptoms, they were given the corresponding symptom code and the earlier date. A first symptom was thus identified for each participant. Many participants reported more than one first symptom, termed 'multiple first symptoms'. 'Subsequent symptoms' were defined as any symptom occurring after a first symptom and prior to diagnosis.

The total diagnostic interval

The **total diagnostic interval (TDI)**, or 'time to diagnosis', was defined as the time from onset of the first symptom/s to the date of diagnosis. Where the date of first presentation to healthcare was known, the **Patient Interval (PI)**, defined as the time from first symptom/s to the first presentation, and the **Health System Interval (HSI)**, defined as the time from first presentation to diagnosis, were also calculated.

Analysis

Descriptive analyses were performed on all demographic, clinical and symptom data for the group as a whole, and by diagnostic group (CRC, all cancer (CRC plus OC), NC). The CRC group was also sub-analysed by stage at diagnosis. The TDI and PI for each group was compared using Wilcoxon rank-sum test.

Clinically relevant demographics, comorbidities, first symptom/s, and family history of cancer variables were *a priori* included in multivariate analyses to identify predictors of time to diagnosis. Variables present in fewer than ten participants were excluded. We chose to analyse CRC, OC and NC groups together for all outcomes because at the time of presentation and referral the final diagnosis was unknown. Flexible parametric survival

models were used to model time to diagnosis, the PI and the HSI. In these cases, rather than death, the event considered to be “failure” in the survival model is either first presentation to healthcare or diagnosis as appropriate. We preferred flexible parametric survival models (Lambert, 2009) over the Cox model as they allow the investigation of duration-varying effects. We present results from two models: firstly, with only time-constant effects and secondly, one that allows for duration-varying effects for symptoms. The former model provides average hazard ratios over time for all variables similar to a Cox model and makes it easier to compare the effects of different factors. The second model allows us to examine in more detail how their effect varies with duration. Symptoms with duration-varying effects were identified using a forward selection approach described in Royston & Lambert (Royston, 2011) and using a significance level of $p \leq 0.01$. All models used splines with five degrees of freedom for the baseline hazard; the duration-dependent effects were modelled using two degrees of freedom for the TDI and HSIs and three degrees of freedom for the PIs.

Sample size

The expected total number of cases in the two areas was 265 annually. Assuming half of these could be identified prospectively from fast track and routine referrals including via colonoscopy (Barrett and Hamilton, 2008), we needed to recruit for 2 years to achieve 200 participants with CRC. We estimated one patient in 25 would have CRC (6-11% in 2-week wait clinics (Rai and Kelly, 2007), but lower out with these clinics). With 200 participants with CRC and at least 10 times as many without we would have over 80% power to detect and $OR > 1.52$ for a common symptom occurring in half of participants. With the same 2200 patients all of which having a TDI we would expect to have 80% power to detect a $HR > 1.23$ for a symptom occurring in 10% of patients.

RESULTS

12,236 patients were approached and 2,667 responded, an overall 21.8% response rate (East 26.3%, North East 18.6%). Of these, 70 did not meet the eligibility criteria, and a further 90 had insufficient data for analysis, leaving a final cohort of 2,507 participants. The demographics of the responders were similar to those of non-responders (responders 47% male, median age 65; non-responders 45% male, median age 65). The stage distribution for those of our cohort who had CRC was comparable to 2013 data for England (late stage 50% vs early 40% and 10% unknown) (NCIN).

The characteristics of the whole cohort are provided in Tables 1 and 2. 152 (6.1%) participants were diagnosed with CRC, 50 (2.0%) with other cancers, and 2,305 (91.9%) with no cancer. The majority of those with CRC had late stage disease ($n=85$, 55.9%; early stage: $n=65$, 42.8%); two CRCs (1.3%) were unstaged. The

proportion of males was higher in the group diagnosed with CRC compared to the group diagnosed with no cancer (57.2% vs 46.2%, $p=0.008$). Those in the CRC group also had a higher median age (71 vs 65, $p<0.001$), and were more likely to have been referred through an urgent pathway (90.1% vs 71.1%, $p<0.001$). The diagnostic groups were otherwise similar in terms of deprivation, education and ethnicity (Table 2).

Among the total cohort, over half had a solitary first symptom (1332, 53.1% (CRC 92, 60.5%, NC 1218, 52.8%). However, multiple first symptoms were common with 21.6% having two first symptoms, 9.3% having three, and 8.4% with four or more multiple first symptoms. 7.6% reported no symptoms first appearing within 2 years of diagnosis (most had symptom pre-dating the 2 year cut-off). Across the total cohort, change in bowel habit (62.8%) and 'bleeding from back passage' (37.8%) were the most common symptoms, but only 45% and 24.4% of those respectively were reported as first symptoms (Table 3). Symptoms not specifically enquired for in the questionnaire but volunteered, such as acute gastro-intestinal illness, perianal pain, flatulence, bloating, and mucus discharge, were individually reported by fewer than 3% of participants. 'Bleeding from back passage' was significantly more frequently reported in people diagnosed with CRC than non-cancer as a first symptom (34.2% vs 23.9%, $p=0.004$). No other symptom as a first or subsequent symptom was more common in those with CRC than those without.

A TDI could be calculated for 2,316 participants; the median TDI across the whole cohort was 136 days (IQR 74-255) (Table 4). There was no difference in the TDI between people diagnosed with CRC and no cancer (CC 124 vs NC 138 days, $p=0.142$, and no evidence of differences between PI and HSI among those with and without CRC (PI median 41 days vs 36 days, $p=0.606$; HSI 49 days vs 59 days, $p=0.078$).

Results from the time constant survival models are shown in Table 5 (supplementary Table A1 for univariate analyses). Older age at diagnosis and a number of symptoms were all associated with a shorter TDI while GI comorbidity, depression/anxiety, and family history of cancer were all associated with a longer TDI and HSI.

Symptom duration-dependent effects

We found symptom duration-dependent effects for 'bleeding from back passage' and change in bowel habit (supplementary Figure 1). The time-constant effects from other variables in the same model are similar to those described in the previous section and are not detailed here. The symptom 'bleeding from back passage' was associated with briefer PI and HSIs initially ($HR>1$), but as the duration of this symptom lengthened the association weakened (HR became close to 1). In other words, when this symptom was present it was associated with faster action (presentation to healthcare professional and/or referral) in the initial period following symptom onset or presentation, compared to other symptoms, but if initial action was not taken the existence of the symptom had little impact on action at later times. In contrast, a change in bowel habit was associated with a lengthier PI initially ($HR<1$) but if the symptom persisted beyond 10 days it became associated with a briefer PI ($HR>1$). In other words, those whose initial symptom was a change in bowel habit

were less likely to present early than those with other symptoms, but of those who did not present in the first 10 days, they presented faster on average than people with other symptoms (HR>1). Change in bowel habit was always associated with a briefer HSI, with no time-dependent effect.

For early stage CRC, the median TDI for any first symptom was 157 days compared with 99 days for late stage CRC ($p=0.019$) (Table 6).

DISCUSSION

This is the first study worldwide to recruit a large prospective cohort of patients with suspicious symptoms before diagnosis in order to study factors associated with time to diagnosis for CRC. Among these patients, who had been referred for investigation, rectal bleeding was the only symptom more frequently seen with CRC, but it occurred in only a third of cases as a first symptom and less than two thirds at any point prior to diagnosis. Other so-called 'alarm' symptoms such as change in bowel habit were also very common in this referred population but did not differentiate between CRC and those without cancer. The positive predictive value of these symptoms will be higher in this study population than in primary care, but the odds ratio between those with and without cancer will fall as this is an enriched, referred population. This doesn't mean these symptoms should not be taken seriously. Multiple first symptoms were common, and symptoms often evolved over time before patients sought health care. There were different symptom and patient factors associated with the PI and HSIs. Some less specific symptoms such as indigestion/abdominal pain or 'feeling different' were associated with shorter PIs; in contrast, only the very specific, 'classical', symptoms such as change in bowel habit and rectal bleeding were associated with shorter HSIs.

Our novel analysis of duration-dependent effects of symptoms showed different patient and healthcare provider responses to certain key symptoms of CRC according to how long the patient had, compared to other symptoms, experienced them. A short duration of rectal bleeding for some patients triggers an early consultation and prompt response by the health care system. Once rectal bleeding has been present for a longer time it is no more likely than other symptoms to have a shorter diagnostic interval despite its recognition as a classical alarm symptom. This means that both patients and healthcare providers may normalise longer-term rectal bleeding and not consult or investigate promptly (Emery et al., 2013).

Comorbidities in the whole cohort were also associated with longer total diagnostic and healthcare intervals. Importantly, people with mental health problems, self-reported anxiety or depression, experienced a longer TDI and HSI, suggesting that their possible physical symptoms were not taken as seriously and were investigated later. Similarly, gastro-intestinal comorbidity increased time to diagnosis, probably due to health care providers attributing new or worsening symptoms to pre-existing illness (Emery et al., 2013, N Hall, 2015).

There was no evidence that those with CRC were diagnosed more quickly than those with an alternative diagnosis. This is surprising given the fact that there were a higher proportion of cancers utilising the urgent pathway, and guidance on urgent referral for suspected CRC in England recommends that 'alarm' symptoms such as rectal bleeding and change in bowel habit warrant urgent referral for investigation (NICE, 2005). Symptoms other than rectal bleeding in this relatively large prospective cohort did not help differentiate CRC from other diagnoses.

The median PI was 35 days and median HSI (combining primary and secondary care) was 58 days. This contrasts with findings of the secondary analysis of CRC data in the National Audit of Cancer Diagnosis in Primary Care, where data from primary care medical records showed a median PI of 19 days (Lyratzopoulos et al., 2015b). In a separate study utilising the same dataset, 21% of CRC patients had 3 or more primary care consultations prior to referral, potentially contributing to longer HSIs (Lyratzopoulos et al., 2013). However, in an analysis of UK General Practice Research Database data for 2962 patients with CRC in 2007/8, the median HSI, of 80 days, was more comparable to our own findings (Neal et al., 2014a).

Key strengths of this study were the prospective design and the collection of data from several sources: patient reports, and primary care and specialist records. The analytical and reporting approaches were robust and performed according to the methodological approaches and definitions recommended in the Aarhus statement (Weller et al., 2012) and STROBE guidelines (von Elm et al., 2007). We chose to define the date of first symptom/s using the patient-reported date rather than the primary care-reported date, as we were analysing patient-reported symptom/s. Ideally, a study would recruit patients from primary care before referral; however, this would be accompanied by major logistical and resource implications in identifying an extremely large prospective cohort of patients with colorectal symptoms in order to capture sufficient patients with cancer. Instead, we recruited patients when they first encountered secondary care; this had the added benefit of allowing us to recruit patients presenting as emergencies, and those referred from other specialists. Recruitment involved two regions of England, selected to ensure a broad range of socio-economic, educational and occupational levels, and the deprivation data suggest that the cohort was reasonably representative of the national population.

The main study limitation is the overall recruitment rate of 22%, though this is similar to other recent studies (Kidney et al., 2015, Walter et al., 2015). We sought to make contact with the target population before they underwent investigation and received a diagnosis, and it is possible that some were unable or unwilling to complete a questionnaire at this worrying time. We are also likely to have under-recruited people who presented as an emergency or who died soon after presentation. However, a recent study from the English Cancer Patient Experience Study showed that only 6% of eligible patients died between sampling and mail-out suggesting potential survival bias in these types of study is relatively small (Abel et al., 2016). The

demographics of non-responders were very similar to participants, and the proportion of late stage CRC was comparable to national data, suggesting our cohort was reasonably representative, that selection bias was not a major issue, and that the findings can be generalised to similar symptomatic populations. If sicker patients were less likely to take part in the study, and they were more likely to have shorter intervals, we may have overestimated the typical intervals in the population. We may well have also underestimated differences between those with different presenting symptoms. In common with a Danish prospective population-based study of diagnostic intervals, the problem of confounding by indication remains to some extent (Torrington et al., 2012). Our results did not find that shorter time to diagnosis was associated with earlier stage disease. It may be that late stage disease has different symptom profiles which affect help-seeking and use of diagnostic pathways. Tumour factors (such as histological type, rate of growth and location) and host factors (such as comorbidity) can influence diagnostic intervals and result in apparently earlier diagnosis of later stage disease. As our study set out to investigate the diagnostic intervals from perception of first symptom to diagnosis, our primary exposure was the initial symptom or symptoms. While this approach allows us to make robust comparisons with other findings, it may obscure the finer detail of symptom patterns and clusters as they evolve over time, and their effects on timely help-seeking by patients and timely diagnosis in primary and secondary care. Although this was a large cohort, we had insufficient power to examine specific clusters of symptoms and their associations with outcomes; a much larger prospective study would be required to achieve this. The fact that fewer CRC cases were recruited than was the aim meant that the study was only powered to detect large differences in symptoms between those with and without CRC.

This study shows that there are subtle differences in the impact of symptoms and patient factors on patient and healthcare intervals, with some clear implications for policy makers and clinicians. Although rectal bleeding was the only symptom predictive of CRC in this referred population it was only reported as the first symptom in one third of cases and as a subsequent symptom in a further 25% of cases. Despite conducting such a large prospective cohort study we failed to identify any other strong solitary symptom signals of CRC, suggesting that bowel cancer awareness campaigns which currently concentrate on a single symptom should also consider messages that reflect the importance of multiple symptoms and evolution of symptoms over time (Moffat et al., 2015). The recently revised NICE guidelines for early detection of CRC support this premise, and have also lowered the threshold for referral, in line with patient preferences for investigation (Banks et al., 2014a, Banks et al., 2014b). However, our study has also shown that only people presenting with shorter histories of rectal bleeding are investigated promptly, and that health care professionals should remain alert to symptoms of possible CRC in people with a history of gastro-intestinal or mental health conditions. The increasingly widespread use of clinical decision support in primary care can also be informed by our findings (Dikomitis et al., 2015, Green et al., 2015), but further research is needed, alongside GPs and

specialists, to identify mechanisms by which patients can be identified, referred and diagnosed in the most timely and appropriate way.

In conclusion, as efforts to expedite the diagnosis of symptomatic colorectal cancer are likely to have benefits for patients in terms of improved survival, earlier-stage diagnosis and improved quality of life, it continues to be a priority to identify symptoms and other factors which should prompt an individual to seek help or a GP to refer in an appropriate and timely manner. It is also important to develop other strategies for earlier diagnosis, including increasing uptake of CRC screening and perhaps the development of biomarkers to improve early detection. Nevertheless, these data provide support for more targeted evaluation of suspicious symptoms in an attempt to identify CRC at an earlier and more amenable stage. It may also be that targeted interventions at higher risk populations aimed at symptom monitoring could be more effective at recognising symptom evolution.

Contributors

FW, JE, GR and WH designed the study. HM oversaw the study's conduct and management, and NH, KM and CD were responsible for conducting and completing data collection. SM analysed the data, supported by GA and previous analyses by CB. All the authors contributed to interpretation of the data. FW, GR and JE wrote the first drafts of the paper and all the authors made critical revisions. The authors fulfil the four criteria for authorship in the ICMJE recommendations.

Funder

This paper presents independent research funded by the National Institute for Health Research (NIHR) Programme Grants for Applied Research programme (RP-PG-0608-10045). The views expressed are those of the authors and not necessarily those of the National Health Service, the National Institute for Health Research, or the Department of Health.

Conflicts of interest

From March 2012 to March 2014 GR was the Royal College of General Practitioners Clinical Lead for Cancer and was a national advocate for the role of the GP in cancer diagnosis. The remaining authors have no conflicts of interest relevant to this research paper.

Acknowledgements

The study team would like to thank all the patients who contributed to this study, and the public representatives in the study team for their valuable contributions to the project: Sue Ballard, Victor Boulter and Margaret Johnson.

We would also like to thank the clinical teams, research nurses and clinic administrators at the participating hospitals: Cambridge University Hospitals NHS Foundation Trust- Mr Miller, Dr Cameron, Kathy Restarick, Debbie Hugh; North Tees & Hartlepool NHS Foundation Trust- Dr Rutter, Sue Kelsey, Debbie Whetter; County Durham and Darlington NHS Foundation Trust-Dr Cundell, Jill Deane, Gill Horner.

Finally, we would like to acknowledge the contribution to the research presented in this paper made by members of the Discovery Programme Steering Committee: Roger Jones (chair); Jon Banks; Alison Clutterbuck; Ardiana Gjini; Joanne Hartland; Maire Justice; Jenny Knowles; Richard Neal; Peter Rose.

Appendix A Inclusion-exclusion recruitment criteria

(i) Fast/urgent clinic track referrals and emergency admissions: include ALL.

(ii) Routine referrals (and via colonoscopy): Include patients with

- Change in bowel habit (including diarrhoea or constipation)
- Rectal bleeding/ blood in stool
- Abdominal pain (in presence of weight loss)
- Abdominal mass
- Anaemia
- Expressed concern about possibility of cancer

Exclude:

- Review of established diagnosis where there is no cancer concern
- Previous GI cancer
- Known infective diarrhoea (e.g. after recent foreign travel)
- Abdominal pain in the absence of weight loss

Appendix B

The questionnaire starts with: *'We are interested in symptoms which you think are related to your recent referral to hospital. The following questions are about when you first noticed a symptom and when you first told your GP or nurse about it. Please give an exact date if you can. Otherwise please give your best estimate (for example, approximately how long ago, the month or the season). You may wish to refer to your diary or calendar if you have it with you'*. The first question was: 'What was the first thing or symptom you noticed that made you think something might be wrong?' Eight specific symptoms followed: change in bowel habit; 'bleeding from the back passage'; back pain; indigestion or abdominal pain 'that wasn't normal for you'; plus decrease in appetite, unexplained weight loss, fatigue or tiredness 'that is unusual for you', and feeling different 'in yourself' from usual.

REFERENCES

- Index of Multiple Deprivation* [Online]. Available: <http://data.gov.uk/dataset/index-of-multiple-deprivation> [Accessed 25th March 2016].
- ABEL, G. A., SAUNDERS, C. L. & LYRATZOPOULOS, G. 2016. Post-sampling mortality and non-response patterns in the English Cancer Patient Experience Survey: Implications for epidemiological studies based on surveys of cancer patients. *Cancer Epidemiol*, 41, 34-41.
- ALLEMANI, C., WEIR, H. K., CARREIRA, H., HAREWOOD, R., SPIKA, D., WANG, X. S., BANNON, F., AHN, J. V., JOHNSON, C. J., BONAVENTURE, A., MARCOS-GRAGERA, R., STILLER, C., AZEVEDO E SILVA, G., CHEN, W. Q., OGUNBIYI, O. J., RACHET, B., SOEBERG, M. J., YOU, H., MATSUDA, T., BIELSKA-LASOTA, M., STORM, H., TUCKER, T. C., COLEMAN, M. P. 2015. Global surveillance of cancer survival 1995-2009: analysis of individual data for 25,676,887 patients from 279 population-based registries in 67 countries (CONCORD-2). *Lancet*, 385, 977-1010.
- BANKS, J., HOLLINGHURST, S., BIGWOOD, L., PETERS, T. J., WALTER, F. M. & HAMILTON, W. 2014a. Preferences for cancer investigation: a vignette-based study of primary-care attendees. *Lancet Oncol*, 15, 232-40.
- BANKS, J., WALTER, F. M., HALL, N., MILLS, K., HAMILTON, W. & TURNER, K. M. 2014b. Decision making and referral from primary care for possible lung and colorectal cancer: a qualitative study of patients' experiences. *Br J Gen Pract*, 64, e775-82.
- BARRETT, J. & HAMILTON, W. 2008. Pathways to the diagnosis of lung cancer in the UK: a cohort study. *BMC Fam. Pract.*, 9, 31.
- BEEKEN, R. J., SIMON, A. E., VON WAGNER, C., WHITAKER, K. L. & WARDLE, J. 2011. Cancer fatalism: deterring early presentation and increasing social inequalities? *Cancer Epidemiol Biomarkers Prev*, 20, 2127-31.
- BISWAS, M., ADES, A. E. & HAMILTON, W. 2015. Symptom lead times in lung and colorectal cancers: what are the benefits of symptom-based approaches to early diagnosis? *Br J Cancer*, 112, 271-7.
- CANCER RESEARCH UK. Available: <http://info.cancerresearchuk.org/cancerstats/mortality/cancerdeaths/> [Accessed 26th March 2016].
- COLEMAN, M. P., FORMAN, D., BRYANT, H., BUTLER, J., RACHET, B., MARINGE, C., NUR, U., TRACEY, E., COORY, M., HATCHER, J., MCGAHAN, C. E., TURNER, D., MARRETT, L., GJERSTORFF, M. L., JOHANNESSEN, T. B., ADOLFSSON, J., LAMBE, M., LAWRENCE, G., MEECHAN, D., MORRIS, E. J.,

- MIDDLETON, R., STEWARD, J. & RICHARDS, M. A. 2011. Cancer survival in Australia, Canada, Denmark, Norway, Sweden, and the UK, 1995-2007 (the International Cancer Benchmarking Partnership): an analysis of population-based cancer registry data. *Lancet*, 377, 127-38.
- DIKOMITIS, L., GREEN, T. & MACLEOD, U. 2015. Embedding electronic decision-support tools for suspected cancer in primary care: a qualitative study of GPs' experiences. *Prim Health Care Res Dev*, 1-8.
- DIN, N. U., UKOUMUNNE, O. C., RUBIN, G., HAMILTON, W., CARTER, B., STAPLEY, S. & NEAL, R. D. 2015. Age and Gender Variations in Cancer Diagnostic Intervals in 15 Cancers: Analysis of Data from the UK Clinical Practice Research Datalink. *PLoS One*, 10, e0127717.
- DREGAN, A., MOLLER, H., CHARLTON, J. & GULLIFORD, M. C. 2013. Are alarm symptoms predictive of cancer survival?: population-based cohort study. *Br J Gen Pract*, 63, e807-12.
- EMERY, J. D., WALTER, F. M., GRAY, V., SINCLAIR, C., HOWTING, D., BULSARA, M., BULSARA, C., WEBSTER, A., AURET, K., SAUNDERS, C., NOWAK, A. & HOLMAN, C. D. 2013. Diagnosing cancer in the bush: a mixed-methods study of symptom appraisal and help-seeking behaviour in people with cancer from rural Western Australia. *Fam Pract*, 30, 294-301.
- ESTEVA, M., LEIVA, A., RAMOS, M., PITA-FERNANDEZ, S., GONZALEZ-LUJAN, L., CASAMITJANA, M., SANCHEZ, M. A., PERTEGA-DIAZ, S., RUIZ, A., GONZALEZ-SANTAMARIA, P., MARTIN-RABADAN, M., COSTA-ALCARAZ, A. M., ESPI, A., MACIA, F., SEGURA, J. M., LAFITA, S., ARNAL-MONREAL, F., AMENGUAL, I., BOSCA-WATTS, M. M., HOSPITAL, A., MANZANO, H. & MAGALLON, R. 2013. Factors related with symptom duration until diagnosis and treatment of symptomatic colorectal cancer. *BMC Cancer*, 13, 87.
- FORBES, L. J., SIMON, A. E., WARBURTON, F., BONIFACE, D., BRAIN, K. E., DESSAIX, A., DONNELLY, C., HAYNES, K., HVIDBERG, L., LAGERLUND, M., LOCKWOOD, G., TISHELMAN, C., VEDSTED, P., VIGMOSTAD, M. N., RAMIREZ, A. J., WARDLE, J. & INTERNATIONAL CANCER BENCHMARKING PARTNERSHIP MODULE 2 WORKING, G. 2013. Differences in cancer awareness and beliefs between Australia, Canada, Denmark, Norway, Sweden and the UK (the International Cancer Benchmarking Partnership): do they contribute to differences in cancer survival? *Br J Cancer*, 108, 292-300.
- GREEN, T., MARTINS, T., HAMILTON, W., RUBIN, G., ELLIOTT, K. & MACLEOD, U. 2015. Exploring GPs' experiences of using diagnostic tools for cancer: a qualitative study in primary care. *Fam Pract*, 32, 101-5.

- HAMILTON, W., ROUND, A., SHARP, D. & PETERS, T. J. 2005. Clinical features of colorectal cancer before diagnosis: a population-based case-control study. *Br.J.Cancer*, 93, 399-405.
- HANSEN, P. L., HJERTHOLM, P. & VEDSTED, P. 2015. Increased diagnostic activity in general practice during the year preceding colorectal cancer diagnosis. *Int J Cancer*, 137, 615-24.
- KEEBLE, S., ABEL, G. A., SAUNDERS, C. L., MCPHAIL, S., WALTER, F. M., NEAL, R. D., RUBIN, G. P. & LYRATZOPOULOS, G. 2014. Variation in promptness of presentation among 10,297 patients subsequently diagnosed with one of 18 cancers: Evidence from a national audit of cancer diagnosis in primary care. *Int J Cancer*, 135(5):1220-8.
- KIDNEY, E., BERKMAN, L., MACHERIANAKIS, A., MORTON, D., DOWSWELL, G., HAMILTON, W., RYAN, R., AWBERY, H., GREENFIELD, S. & MARSHALL, T. 2015. Preliminary results of a feasibility study of the use of information technology for identification of suspected colorectal cancer in primary care: the CREDIBLE study. *Br J Cancer*, 112 Suppl 1, S70-6.
- LAMBERT, P. C. Rotston, P. 2009. Further development of flexible parametric models for survival analysis. *Stata Journal*, 9, 265-290.
- LYRATZOPOULOS, G., ABEL, G. A., MCPHAIL, S., NEAL, R. D. & RUBIN, G. P. 2013. Measures of promptness of cancer diagnosis in primary care: secondary analysis of national audit data on patients with 18 common and rarer cancers. *Br J Cancer*, 108, 686-90.
- LYRATZOPOULOS, G., LIU, M. P., ABEL, G. A., WARDLE, J. & KEATING, N. L. 2015a. The Association between Fatalistic Beliefs and Late Stage at Diagnosis of Lung and Colorectal Cancer. *Cancer Epidemiol Biomarkers Prev*, 24, 720-6.
- LYRATZOPOULOS, G., SAUNDERS, C. L., ABEL, G. A., MCPHAIL, S., NEAL, R. D., WARDLE, J. & RUBIN, G. P. 2015b. The relative length of the patient and the primary care interval in patients with 28 common and rarer cancers. *Br J Cancer*, 112 Suppl 1, S35-40.
- MACLEOD, U., MITCHELL, E. D., BURGESS, C., MACDONALD, S. & RAMIREZ, A. J. 2009. Risk factors for delayed presentation and referral of symptomatic cancer: evidence for common cancers. *Brit J Cancer*, 101 Suppl 2, S92-S101.
- MARINGE, C., WALTERS, S., RACHET, B., BUTLER, J., FIELDS, T., FINAN, P., MAXWELL, R., NEDREBO, B., PAHLMAN, L., SJOVALL, A., SPIGELMAN, A., ENGHOLM, G., GAVIN, A., GJERSTORFF, M. L., HATCHER, J., JOHANNESSEN, T. B., MORRIS, E., MCGAHAN, C. E., TRACEY, E., TURNER, D., RICHARDS, M. A.,

- COLEMAN, M. P. 2013. Stage at diagnosis and colorectal cancer survival in six high-income countries: a population-based study of patients diagnosed during 2000-2007. *Acta Oncol*, 52, 919-32.
- MCLACHLAN, S., MANSELL, G., SANDERS, T., YARDLEY, S., VAN DER WINDT, D., BRINDLE, L., CHEW-GRAHAM, C. & LITTLE, P. 2015. Symptom perceptions and help-seeking behaviour prior to lung and colorectal cancer diagnoses: a qualitative study. *Fam Pract*, 32(5):568-77.
- MOFFAT, J., BENTLEY, A., IRONMONGER, L., BOUGHEY, A., RADFORD, G. & DUFFY, S. 2015. The impact of national cancer awareness campaigns for bowel and lung cancer symptoms on sociodemographic inequalities in immediate key symptom awareness and GP attendances. *Br J Cancer*, 112 Suppl 1, S14-21.
- N HALL, L. B., J BANKS, J EMERY, K MILLS, M JOHNSON, GP RUBIN, W HAMILTON, FM WALTER 2015. Symptom appraisal and healthcare-seeking for symptoms suggestive of colorectal cancer: a qualitative study. *BMJ Open*, 5(10):e008448.
- National Cancer Intelligence Network (NCIN).
http://www.ncin.org.uk/cancer_type_and_topic_specific_work/cancer_type_specific_work/colorectal_cancer/ [Accessed 25th March 2016].
- NEAL, R. D. & ALLGAR, V. L. 2005. Sociodemographic factors and delays in the diagnosis of six cancers: analysis of data from the "National Survey of NHS Patients: Cancer". *Br J Cancer*, 92, 1971-1975.
- NEAL, R. D., DIN, N. U., HAMILTON, W., UKOUMUNNE, O. C., CARTER, B., STAPLEY, S. & RUBIN, G. 2014a. Comparison of cancer diagnostic intervals before and after implementation of NICE guidelines: analysis of data from the UK General Practice Research Database. *Br J Cancer*, 110, 584-92.
- NEAL, R. D., NAFEES, S., PASTERFIELD, D., HOOD, K., HENDRY, M., GOLLINS, S., MAKIN, M., STUART, N., TURNER, J., CARTER, B., WILKINSON, C., WILLIAMS, N. & ROBLING, M. 2014b. Patient-reported measurement of time to diagnosis in cancer: development of the Cancer Symptom Interval Measure (C-SIM) and randomised controlled trial of method of delivery. *BMC Health Serv Res*, 14, 3.
- NEAL, R. D., THARMANATHAN, P., FRANCE, B., DIN, N. U., COTTON, S., FALLON-FERGUSON, J., HAMILTON, W., HENDRY, A., HENDRY, M., LEWIS, R., MACLEOD, U., MITCHELL, E. D., PICKETT, M., RAI, T., SHAW, K., STUART, N., TORRING, M. L., WILKINSON, C., WILLIAMS, B., WILLIAMS, N. & EMERY, J. 2015. Is increased time to diagnosis and treatment in symptomatic cancer associated with poorer outcomes? Systematic review. *Br J Cancer*, 112 Suppl, S92-S107.

- NICE. 2005. *Referral guidelines for suspected cancer (CG 27)* [Online]. Available: <http://www.nice.org.uk/CG027> [Accessed 25th March 2016].
- QUAIFE, S. L., FORBES, L. J., RAMIREZ, A. J., BRAIN, K. E., DONNELLY, C., SIMON, A. E. & WARDLE, J. 2014. Recognition of cancer warning signs and anticipated delay in help-seeking in a population sample of adults in the UK. *Br J Cancer*, 110, 12-8.
- RAI, S. & KELLY, M. J. 2007. Prioritization of colorectal referrals: a review of the 2-week wait referral system. *Colorectal Dis*, 9, 195-202.
- RASMUSSEN, S., LARSEN, P. V., SONDERGAARD, J., ELNEGAARD, S., SVENDSEN, R. P. & JARBOL, D. E. 2015. Specific and non-specific symptoms of colorectal cancer and contact to general practice. *Fam Pract*, 32, 387-94.
- ROYSTON, P. Lambert, P.C. 2011. *Flexible parametric survival analysis using Stata: beyond the Cox model*. College Station, TX, Stata Press., TX, Stata Press.
- RUBIN, G., WALTER, F.M, EMERY, J., NEAL, R., HAMILTON, W. & WARDLE, J. 2014. Research into practice: prompt diagnosis of cancer in primary care. *Br J Gen Pract*, 64, 428-30.
- SIMON, A. E., WALLER, J., ROBB, K. & WARDLE, J. 2010. Patient delay in presentation of possible cancer symptoms: the contribution of knowledge and attitudes in a population sample from the United Kingdom. *Cancer Epidemiol Biomarkers Prev*, 19, 2272-7.
- TORRING, M. L., FRYDENBERG, M., HAMILTON, W., HANSEN, R. P., LAUTRUP, M. D. & VEDSTED, P. 2012. Diagnostic interval and mortality in colorectal cancer: U-shaped association demonstrated for three different datasets. *J Clin Epidemiol*, 65, 669-78.
- TORRING, M. L., FRYDENBERG, M., HANSEN, R. P., OLESEN, F., HAMILTON, W. & VEDSTED, P. 2011. Time to diagnosis and mortality in colorectal cancer: a cohort study in primary care. *Brit J Cancer*, 104, 934-40.
- TRAVIS, S. P., SCHNELL, D., KRZESKI, P., ABREU, M. T., ALTMAN, D. G., COLOMBEL, J. F., FEAGAN, B. G., HANAUER, S. B., LEMANN, M., LICHTENSTEIN, G. R., MARTEAU, P. R., REINISCH, W., SANDS, B. E., YACYSHYN, B. R., BERNHARDT, C. A., MARY, J. Y. & SANDBORN, W. J. 2012. Developing an instrument to assess the endoscopic severity of ulcerative colitis: the Ulcerative Colitis Endoscopic Index of Severity (UCEIS). *Gut*, 61(4):535-42.

- VON ELM, E., ALTMAN, D. G., EGGER, M., POCOCK, S. J., GOTZSCHE, P. C., VANDENBROUCKE, J. P. & INITIATIVE, S. 2007. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Brit Med J*, 335, 806-8.
- WALTER, F. M., RUBIN, G., BANKHEAD, C., MORRIS, H. C., HALL, N., MILLS, K., DOBSON, C., RINTOUL, R. C., HAMILTON, W. & EMERY, J. 2015. Symptoms and other factors associated with time to diagnosis and stage of lung cancer: a prospective cohort study. *Br J Cancer*, 112 Suppl, S6-S13.
- WELLER, D., VEDSTED, P., RUBIN, G., WALTER, F. M., EMERY, J., SCOTT, S., CAMPBELL, C., ANDERSEN, R. S., HAMILTON, W., OLESEN, F., ROSE, P., NAFEES, S., VAN RIJSWIJK, E., HIOM, S., MUTH, C., BEYER, M. & NEAL, R. D. 2012. The Aarhus statement: improving design and reporting of studies on early cancer diagnosis. *Br J Cancer*, 106, 1262-7.

Table 1: Diagnostic characteristics of participants

	Total (%)
Primary Colorectal Cancer	152 (6.1)
Other Cancers:	50 (2.0)
Lymphoma	9
Prostate cancer	7
Gynaecological cancer (ovarian, uterine)	7
Anal cancer	5
Renal cancer (kidney, bladder)	4
CUP	4
Oesophageal cancer	3
Pancreatic cancer	2
Multiple myeloma	2
Stomach cancer	2
Other ¹	5
Non Cancer ¹ :	2305 (91.9)
Polyps	612 (26.6)
Polyp with metaplasia	200 (8.7)
Poly with dysplasia	168 (7.3)
Polyp NOS	244 (10.6)
Nil abnormal	598 (25.9)
Diverticular disease	463 (20.1)
Inflammatory Bowel Disease (IBD)	189 (8.2)
Colitis ulcerative	55 (2.4)
Colitis - other	99 (4.3)
Crohn's Disease	7 (0.3)
IBD NOS	28 (1.2)
Haemorrhoids	186 (8.1)
Irritable Bowel Syndrome	95 (4.1)
Benign gastro-duodenal disease	83 (3.6)
Miscellaneous	51 (2.2)
Anaemia (no lower GI cause found)	31 (1.3)
Hepato-pancreato-biliary disorders	20 (0.9)
Anal conditions (excluding haemorrhoids)	13 (0.6)
Constipation	13 (0.6)
Coeliac disease	5 (0.2)
Total	2507

Notes:

1) Percentages add to more than 100% because 54 patients have more than one diagnosis. More than one diagnosis in the same category (e.g. miscellaneous) was counted only once.

2) Other cancers include (n=1): leiomyosarcoma, appendix, peritoneal, melanoma, cholangiocarcinoma, CUP=Cancer Unknown Primary, including one diagnosed as 'metastatic', NOS=not otherwise specified

Table 2: Socio-demographic characteristics of participants

Values are n (%) unless otherwise stated

	Colorectal Cancer (CRC) (n = 152)	Other Cancer (OC) (n = 50)	No Cancer (NC) (n = 2305)	Total Cohort ¹ (n = 2507)	p-value CRC vs NC
Gender					
Male	87 (57.2)	30 (60)	1064 (46.2)	1181 (47.1)	0.008
Female	65 (42.8)	20 (40)	1241 (53.8)	1326 (52.9)	
Age (median, range)					
	71 (40-92)	71.5 (51-91)	65 (40-100)	65 (40-100)	<0.001
Highest education level					
Degree/Diploma/equivalent	67 (44.1)	17 (34.0)	941 (40.8)	1025 (40.9)	0.511
A Level/GCSE/O Level	42 (27.6)	16 (32.0)	740 (32.1)	798 (31.8)	
Other/None/missing	43 (28.3)	17 (34.0)	624 (27.1)	684 (27.3)	
Ethnicity					
White	149 (98.0)	50 (100)	2259 (98.0)	2458 (98.1)	0.640
Other/missing	3 (2.0)	0	46 (2.0)	49 (2.0)	
Smoking status					
Current	16 (10.5)	2 (4.0)	205 (8.9)	223 (8.9)	0.549
Ex-smoker	68 (44.7)	21 (42.0)	970 (42.1)	1059 (42.2)	
Never/missing	68 (44.7)	27 (54.0)	1130 (49.0)	1225 (48.9)	
Lives alone					
Yes	43 (28.3)	14 (28.0)	538 (23.3)	595 (23.7)	0.164
No/missing	109 (71.8)	36 (72.0)	1767 (76.7)	1912 (76.3)	
Deprivation (IMD- quintiles)					
1 st National Quintile (least)	61 (40.1)	13 (26.0)	826 (35.8)	900 (35.9)	0.168
2 nd National Quintile	40 (26.3)	12 (24.0)	550 (23.9)	602 (24.0)	
3 rd National Quintile	12 (7.9)	10 (20.0)	352 (15.3)	374 (14.9)	
4 th National Quintile	20 (13.2)	10 (20.0)	304 (13.2)	334 (13.3)	
5 th National Quintile (most)	19 (12.5)	5 (10.0)	263 (11.4)	287 (11.4)	
missing	0	0	10 (0.4)	10 (0.4)	
Referral route					
Urgent	137 (90.1)	44 (88.0)	1638 (71.1)	1819 (72.6)	<0.001
GP FT	124 (90.5)	42 (95.5)	1597 (97.5)	1763 (96.9)	
A&E	6 (4.4)	1 (2.3)	14 (0.9)	21 (1.2)	
Other	7 (5.1)	1 (2.3)	27 (1.6)	17 (0.9)	
Non-urgent	15 (9.9)	6 (12.0)	667 (28.9)	688 (27.4)	
Region					
Cambridge	78 (51.3)	21 (42.0)	1109 (48.1)	1208 (48.2)	0.444
North East	74 (48.7)	29 (58.0)	1196 (51.9)	1299 (51.8)	

Missing values were included in the Other/Never categories where possible. Percentages are as follows: Employment Status (2.0), Education Level (3.3), Ethnicity (0.8), Smoking Status (1.5), Live Alone (0.6), referral route (2.2).

Table 3: Symptoms reported by participants, stratified by all and first symptom/s and diagnostic group

Values are n (%) unless otherwise stated, and all columns add up to >100% because of multiple symptoms.

Symptom	Colorectal Cancer (CRC) (n = 152)		No Cancer (NC) (n = 2305)		Total Cohort (n = 2507)		P-value for first symptoms CRC vs NC
	First symptoms	Subsequent symptoms	First symptoms	Subsequent symptoms	First symptoms	Subsequent symptoms	
Symptoms in questionnaire							
Change in bowel habit	66 (43.4)	27 (17.8)	1043 (45.2)	410 (17.8)	1129 (45)	446 (17.8)	0.661
'Bleeding from back passage'	52 (34.2)	39 (25.7)	552 (23.9)	296 (12.8)	612 (24.4)	336 (13.4)	0.004
Indigestion or heartburn or persistent tummy ache that wasn't normal for you	30 (19.7)	13 (8.6)	509 (22.1)	281 (12.2)	549 (21.9)	302 (12)	0.499
Fatigue or tiredness that is unusual for you	35 (23)	23 (15.1)	404 (17.5)	307 (13.3)	448 (17.9)	347 (13.8)	0.086
Feeling different "in yourself" from usual	27 (17.8)	11 (7.2)	391 (17)	257 (11.1)	433 (17.3)	278 (11.1)	0.799
Decrease in appetite	15 (9.9)	16 (10.5)	195 (8.5)	192 (8.3)	220 (8.8)	221 (8.8)	0.547
Back pain	3 (2)	9 (5.9)	197 (8.5)	179 (7.8)	210 (8.4)	194 (7.7)	0.004
Unexplained weight loss	6 (3.9)	18 (11.8)	111 (4.8)	183 (7.9)	127 (5.1)	214 (8.5)	0.626
Other symptoms reported by participants							
Other (non-gastro-intestinal) symptoms	4 (2.6)	3 (2)	47 (2)	16 (0.7)	53 (2.1)	20 (0.8)	NA
Acute gastro-intestinal illness	0 (0)	3 (2)	48 (2.1)	15 (0.7)	49 (2)	18 (0.7)	NA
Perianal pain or discomfort	2 (1.3)	1 (0.7)	37 (1.6)	15 (0.7)	39 (1.6)	16 (0.6)	NA
Wind or flatulence	0 (0)	2 (1.3)	33 (1.4)	11 (0.5)	33 (1.3)	13 (0.5)	NA
Urgency or leakage of bowels	0 (0)	0	31 (1.3)	13 (0.6)	32 (1.3)	13 (0.5)	NA
Bloating	0 (0)	1 (0.7)	26 (1.1)	12 (0.5)	28 (1.1)	14 (0.6)	NA
Mucus or discharge per rectum	0 (0)	2 (1.3)	25 (1.1)	7 (0.3)	25 (1)	9 (0.4)	NA
Pain, non-abdominal	1 (0.7)	0	22 (1)	9 (0.4)	23 (0.9)	9 (0.4)	NA

NA: Not applicable

Table 4: Total Diagnostic Interval, Patient Interval and Health System Interval for first symptom/s for total cohort and by diagnostic groups P-values for CC vs NC with any symptom: (A) TDI p=0.142, (B) PI p=0.606, (C) HSI p=0.078. Similar results were obtained for the comparison between all cancers and NC (data not shown).

Symptom	Total cohort		Colorectal Cancer (CC)		No Cancer (NC)	
	Median (IQR)	n	Median (IQR)	n	Median (IQR)	n
(A) TOTAL DIAGNOSTIC INTERVAL (TDI)*						
Any Symptom	136 (74-255)	2316	124 (74-226)	149	138 (74-264)	2120
Change in bowel habit	113 (69-203)	1129	124 (66-226)	66	112 (70-203)	1043
'Bleeding from back passage'	90 (49-164)	612	88 (47-142)	52	89 (49-173)	552
Indigestion/heartburn/persistent tummy ache	139 (81-277)	549	125 (80-262)	30	142 (81-283)	509
Fatigue or tiredness that is unusual for you	143 (86-239)	448	130 (92-189)	35	146 (86-247)	404
Feeling different "in yourself" from usual	128 (77-216)	433	124 (80-154)	27	130 (76-231)	391
Decrease in appetite	115 (69-235)	220	98 (64-138)	15	115 (69-269)	195
Back pain	161 (79-294)	210	98 (92-525)	3	162 (79-295)	197
Unexplained weight loss	148 (84-333)	127	111 (82-205)	6	159 (86-346)	111
(B) PATIENT INTERVAL (PI)**						
Any Symptom	35 (7-92)	2103	41 (13-92)	128	36 (7-93)	1932
Change in bowel habit	42 (14-92)	1106	57 (22-115)	63	42 (14-92)	1023
'Bleeding from back passage'	25 (3-69)	607	29 (6-66)	52	24 (3-69)	547
Indigestion/heartburn/persistent tummy ache	31 (5-86)	525	28 (7-90)	27	31 (5-90)	488
Fatigue or tiredness that is unusual for you	31 (7-89)	365	35 (8-65)	26	31 (6-90)	331
Feeling different "in yourself" from usual	30 (4-76)	377	37 (14-65)	21	30 (3-77)	343
Decrease in appetite	20 (3-62)	190	33 (8-90)	15	18 (2-62)	166
Back pain	29 (2-76)	183	41 (37-64)	3	29 (1-88)	172
Unexplained weight loss	31 (5-92)	115	46 (35-73)	4	31 (2-96)	102
(C) HEALTH CARE INTERVAL (HCI)**						
Any Symptom	58 (27-128)	2103	49 (26-106)	128	59 (27-129)	1932
Change in bowel habit	51 (25-99)	1106	47 (28-101)	63	52 (25-99)	1023
'Bleeding from back passage'	41 (19-85)	607	37 (20-60)	52	41 (19-86)	547
Indigestion/heartburn/persistent tummy ache	76 (34-156)	525	58 (35-151)	27	78 (34-161)	488
Fatigue or tiredness that is unusual for you	69 (30-146)	365	54 (27-89)	26	70 (29-147)	331
Feeling different "in yourself" from usual	67 (32-139)	377	61 (38-87)	21	69 (30-144)	343
Decrease in appetite	60 (31-138)	190	49 (30-63)	15	64 (30-158)	166
Back pain	88 (35-203)	183	61 (51-461)	3	82 (35-206)	172
Unexplained weight loss	83 (39-191)	115	33 (16-50)	4	89 (43-202)	102

*As these are median time intervals (rather than mean intervals) there is no expectation that the median TDI will equal the sum of median PI and median HSI.

**The PI and HSI could only be calculated for 2,103 participants (unknown presentation date n=213). Those with an available date of presentation had a shorter median TDI (130 days vs 199 days for the remaining cases)

Table 5: Predictors of Total Diagnostic Interval (TDI), Patient Interval (PI) and Health System Interval (HIS)

Hazard ratio estimates are from a flexible parametric survival multivariable model with only time constant effects. Model is adjusted for all variables in table plus ethnicity (non-white vs. white), smoking status (current and ex-smoker vs. never), living alone (yes vs. no) and region (North East vs. Cambridge). **In this context the hazard ratio represents the relative increase in rate of presentation/diagnosis. A hazard ratio of 2 would imply that patients in one group presented/were diagnosed twice as quickly as in the reference group.**

	Total Diagnostic Interval (n=2306)		Patient Interval (n=2095)		Health System Interval (n=2095)	
	HR (95 CI)	p-value	HR (95 CI)	p-value	HR (95 CI)	p-value
Bleeding from Back Passage	1.79 (1.62-1.98)	<0.001	1.41 (1.27-1.57)	<0.001	1.56 (1.41-1.72)	<0.001
Change in Bowel Habit	1.55 (1.42-1.69)	<0.001	1.04 (0.95-1.13)	0.440	1.55 (1.42-1.70)	<0.001
Back pain	0.99 (0.85-1.14)	0.874	1.20 (1.02-1.40)	0.024	0.86 (0.73-1.00)	0.052
Indigestion/Heartburn/Tummy Ache	1.00 (0.90-1.10)	0.942	1.17 (1.06-1.30)	0.003	0.84 (0.76-0.93)	0.001
Decrease in Appetite	1.10 (0.95-1.28)	0.200	1.15 (0.98-1.35)	0.087	1.11 (0.94-1.30)	0.224
Weight loss	0.90 (0.75-1.09)	0.284	1.00 (0.82-1.22)	0.987	0.81 (0.66-0.99)	0.038
Fatigue or Tiredness	1.14 (1.02-1.27)	0.021	1.11 (0.98-1.25)	0.096	1.04 (0.92-1.17)	0.549
Feeling Different	1.21 (1.08-1.35)	0.001	1.25 (1.10-1.41)	<0.001	1.01 (0.90-1.14)	0.812
Gender (ref=Female)						
Male	1.01 (0.93-1.10)	0.739	1.03 (0.94-1.13)	0.535	1.02 (0.93-1.11)	0.731
Age at diagnosis (10 years)	1.08 (1.04-1.12)	<0.001	1.13 (1.08-1.18)	<0.001	0.99 (0.95-1.03)	0.698
IMD (ref=least deprived)		0.611		0.645		0.528
2 nd National Quintile	0.98 (0.88-1.09)		1.03 (0.91-1.15)		0.92 (0.82-1.03)	
3 rd National Quintile	1.03 (0.90-1.17)		1.09 (0.95-1.25)		0.94 (0.82-1.08)	
4 th National Quintile	0.91 (0.79-1.04)		0.97 (0.84-1.12)		0.90 (0.78-1.04)	
5 th National Quintile	0.97 (0.83-1.13)		0.98 (0.83-1.16)		0.96 (0.81-1.13)	
GI Comorbidity	0.80 (0.71-0.89)	<0.001	1.01 (0.89-1.14)	0.904	0.78 (0.69-0.88)	<0.001
Depression/anxiety	0.86 (0.77-0.96)	0.007	1.05 (0.93-1.18)	0.415	0.80 (0.71-0.90)	<0.001
Family History of cancer	0.91 (0.83-0.99)	0.031	0.94 (0.86-1.04)	0.237	0.90 (0.82-0.99)	0.033

Table 6: Time to diagnosis (days) for first symptom/s among colorectal cancer group, stratified by stage at diagnosis

P-values for early vs late stage with any symptom: (i) p=0.019, (ii) p=0.022.

	Early Stage (n = 64)		Late Stage (n = 82)	
	Median (IQR)	n	Median (IQR)	n
Any Symptom	157 (93-243)	64	99 (65-198)	83
Change in bowel habit	159 (92-274)	23	111 (65-173)	41
'Bleeding from back passage'	92 (36-156)	23	84 (47-133)	28
Indigestion or heartburn or persistent tummy ache that wasn't normal for you	188 (96-310)	11	103 (62-153)	18
Fatigue or tiredness that is unusual for you	154 (95-226)	15	110 (81-152)	20
Feeling different "in yourself" from usual	114 (95-154)	6	101 (65-153)	18
Decrease in appetite	138 (95-186)	3	83 (56-128)	11
Back pain	-	0	98 (92-525)	3
Unexplained weight loss	205 (82-572)	3	69 (39-98)	2

^a not staged: 2

SUPPLEMENTARY ONLINE MATERIAL

Table 1: Univariable analysis of factors associated with Total Diagnostic Interval (TDI), Patient Interval (PI) and Health System Interval (HSI)

Hazard ratio estimates are from flexible parametric survival univariate models.

	Total Diagnostic Interval N=2316		Patient Interval N=2103		Health System Interval N=2103	
	HR (95 CI)	p-value	HR (95 CI)	p-value	HR (95 CI)	p-value
'Bleeding from Back Passage'	1.59 (1.45-1.75)	<0.001	1.25 (1.14-1.37)	<0.001	1.52 (1.38-1.67)	<0.001
Change in Bowel Habit	1.43 (1.32-1.55)	<0.001	0.97 (0.89-1.05)	0.424	1.46 (1.34-1.60)	<0.001
Back pain	0.88 (0.76-1.02)	0.081	1.21 (1.04-1.41)	0.015	0.77 (0.66-0.89)	0.001
Indigestion/Heartburn/Tummy Ache	0.92 (0.84-1.01)	0.091	1.13 (1.02-1.24)	0.017	0.78 (0.71-0.86)	<0.001
Decrease in Appetite	1.07 (0.93-1.23)	0.322	1.33 (1.14-1.54)	<0.001	0.99 (0.86-1.15)	0.932
Weight loss	0.83 (0.70-1.00)	0.047	1.03 (0.86-1.25)	0.726	0.79 (0.66-0.95)	0.015
Fatigue or Tiredness	0.99 (0.89-1.10)	0.833	1.13 (1.01-1.27)	0.032	0.91 (0.82-1.02)	0.119
Feeling Different	1.14 (1.02-1.26)	0.017	1.30 (1.16-1.45)	<0.001	0.97 (0.86-1.08)	0.538
Gender (ref=Female)						
Male	1.10 (1.01-1.19)	0.027	1.04 (0.95-1.13)	0.367	1.08 (0.99-1.18)	0.083
Age at diagnosis (10 years)	1.06 (1.02-1.10)	0.001	1.10 (1.06-1.14)	<0.001	1.01 (0.98-1.05)	0.527
Ethnicity (ref= White)						
Other	0.75 (0.55-1.04)	0.083	0.83 (0.59-1.16)	0.270	0.75 (0.53-1.05)	0.090
Smoking Status (ref= Never)		0.702		0.236		0.292
Current	0.94 (0.81-1.10)		1.05 (0.90-1.23)		0.94 (0.81-1.10)	
Ex-Smoker	1.01 (0.92-1.10)		0.94 (0.86-1.03)		1.05 (0.96-1.15)	
Live Alone (ref= No)						
Yes	0.97 (0.88-1.06)	0.468	1.05 (0.95-1.17)	0.306	0.92 (0.83-1.02)	0.098
IMD (ref=least deprived)		0.644		0.334		0.068
2 nd National Quintile	1.00 (0.90-1.12)		1.05 (0.93-1.17)		1.00 (0.89-1.11)	
3 rd National Quintile	1.04 (0.92-1.18)		1.15 (1.01-1.31)		1.02 (0.89-1.16)	
4 th National Quintile	1.04 (0.91-1.18)		1.02 (0.89-1.16)		1.06 (0.93-1.22)	
5 th National Quintile	1.11 (0.96-1.27)		1.06 (0.92-1.23)		1.23 (1.06-1.42)	
GI Comorbidity	0.76 (0.68-0.85)	<0.001	1.05 (0.93-1.18)	0.442	0.73 (0.65-0.82)	<0.001
Depression/anxiety	0.83 (0.75-0.93)	0.001	1.03 (0.92-1.16)	0.560	0.79 (0.70-0.88)	<0.001
Family History of cancer	0.89 (0.82-0.97)	0.011	0.94 (0.85-1.03)	0.157	0.93 (0.84-1.01)	0.098
Region (ref=Cambridge)						
North East	1.33 (1.22-1.44)	<0.001	1.08 (0.99-1.17)	0.093	1.51 (1.38-1.64)	<0.001

SUPPLEMENTARY ONLINE MATERIAL

Figure 1: Time dependent effects of 'bleeding from back passage' and 'change in bowel habit' on the hazard ratio (HR) of Total Diagnostic Interval (TDI), Patient Interval (PI) and Health System Interval (HSI)

